

Critical Areas Report

Newport Way Improvements Project Issaquah, Washington

for **KPG, Inc.**

May 7, 2020



1101 South Fawcett Avenue, Suite 200 Tacoma, Washington 98402 253.383.4940

Critical Areas Report

Newport Way Improvements Project Issaquah, Washington

File No. 0252-039-01

May 7, 2020

Prepared for:

KPG, Inc. 3131 Elliott Avenue, Suite 400 Seattle, Washington 98121

Attention: Sessyle Asato, PE

Prepared by:

GeoEngineers, Inc. 1101 South Fawcett Avenue, Suite 200 Tacoma, Washington 98402 253.383.4940

Emily R. Hurn

Environmental Scientist

David B. Conlin, MS, PWS

Senior Biologist

Joseph O. Callaghan, MS, PWS

Principal Biologist

ERH:DBC:JOC:tlm

Disclaimer: Any electronic form, facsimile or hard copy of the original document (email, text, table, and/or figure), if provided, and any attachments are only a copy of the original document. The original document is stored by GeoEngineers, Inc. and will serve as the official document of record.



Table of Contents

1.0 INTRODUCTION	1
1.1. Project Location	
1.2. Regulatory Requirements	
2.0 METHODS	1
2.1. Data Review	1
2.1.1. Wetland and Stream Data	2
2.1.2. Soil Survey	3
2.2. Literature Review	3
2.2.1. Gateway Apartments - Previous Critical Areas Study and Final Mitigation Plan	3
2.2.2. Riva Townhomes Site Plans	
2.2.3. Former Bergsma Subdivision – Previous Culvert Survey	
2.3. Field Methods	
2.3.1. Wetland Assessment	
2.3.2. Stream Assessment	
2.3.3. Fish and Wildlife Habitat	
3.0 FINDINGS	5
3.1. Wetlands	6
3.2. Stream Delineation	ε
3.3. Stormwater Features	6
3.4. Fish and Wildlife Habitat	7
4.0 SUMMARY	7
5.0 LIMITATIONS	7
6.0 REFERENCES	

LIST OF FIGURES

Figure 1. Vicinity Map

Figure 2. Area of Investigation

Figures 3 through 5. Critical Area Results

APPENDICES

Appendix A. Data Review Sources

Appendix B. Site Photographs

Figures B-1 through B-11. Site Photographs

Appendix C. Sample Plot Data Form

Appendix D. Wetland Rating Form

Figure D-1. Boundaries and Characteristics of Wetland A

Figure D-2. Landscape Characteristics of Wetland A

Figure D-3. Contributing Basin

Figure D-4. 303(d) Screen Shot

Figure D-5. TMDL Screen Shot

Appendix E. Wetland and Stream Descriptions



1.0 INTRODUCTION

GeoEngineers, Inc. (GeoEngineers) was contracted by KPG, Inc. (KPG) to perform wetland and stream delineation services along Newport Way NW between State Route (SR) 900 and SE 54th Street, for the proposed Newport Way Improvements Project (project). The City of Issaquah (City) is planning road improvement actions along the approximate 1.1-mile stretch of road (project corridor). The improvements include, but are not limited to, road widening and culvert replacements. This report has been prepared to provide baseline wetland, stream and wildlife habitat information within the project corridor in accordance with City of Issaquah Municipal Code (IMC), Chapter 18.10 Environmental Protection.

1.1. Project Location

The project is located along Newport Way in Issaquah, Washington (Figure 1, Vicinity Map) and extends from approximately SR 900 to SE 54th Street. The proposed project is within Section 29 of Township 24 North, Range 06 East of the Willamette Meridian (W.M.). Newport Way is a two-lane asphalt roadway. Portions of Newport Way also include sidewalks and bike lane features. The surrounding vicinity largely consists of single- and multi-family residences with some businesses at the south end of the project area. The City is planning road improvement actions along the approximate 1.1-mile stretch of road.

1.2. Regulatory Requirements

This report was prepared to address critical areas review requirements as presented in IMC 18.10, which includes streams, wetlands and associated critical area buffers. The results are also equally applicable for documentation of baseline aquatic habitats necessary for State Environmental Policy Act (SEPA) review, National Environmental Policy Act (NEPA) review, state Hydraulic Project Approval (HPA), and/or federal Clean Water Act permitting requirements.

2.0 METHODS

Methods used to assess wetland, stream and wildlife habitats, occurring within or adjacent to the project corridor, included a review of published data and literature, as well as field assessment activities, as described in more detail in the following sections. The Area of Investigation defined for this effort included up to 100 feet on either side of the road right-of-way (ROW), with the exception of parcels for which right-of-entry (ROE) had not been granted at the time of our fieldwork, as illustrated on Figure 2, Area of Investigation.

2.1. Data Review

GeoEngineers first researched existing information on wetlands and streams documented within the project area vicinity both within and beyond the Area of Investigation. Our search for pertinent and applicable data and maps consisted of a review of the following information sources.

Public Databases:

 Washington Department of Fish and Wildlife (WDFW) Priority Habitat and Species (PHS) database (WDFW 2017a);



- United States Fish and Wildlife Service (USFWS) National Wetlands Inventory (NWI) maps (USFWS 2017a);
- USFWS 2017 List of Threatened and Endangered Species (USFWS 2017b)
- United States Department of Agriculture National Resource Conservation Service (USDA-NRCS)
 Web Soil Survey (USDA-NRCS 2017a);
- Washington State Department of Natural Resources (WDNR) Forest Practices Application Review System (FPARS) (WDNR 2017); and
- WDFW SalmonScape mapping application (WDFW 2017b).

Prior Studies:

- City of Issaquah GIS Data Viewer (City of Issaquah 2017a).
- Gateway Apartments Critical Areas Study and Final Mitigation Plan (Talasaea 2016).
- Bergsma Culvert Survey (CEC 2016; PACE 2016).

Mapped data reports available from WDFW, USFWS, and USDA-NRCS are included in Appendix A, Data Review Sources.

2.1.1. Wetland and Stream Data

The USFWS NWI online mapper (USFWS 2017a) does not show any mapped wetlands within the project area (Appendix A).

The WDNR FPARS map (WDNR 2017) depicts four streams crossing the project corridor: Tibbetts Creek, Anti-Aircraft Creek, Schneider Creek and one unnamed stream depicted as a tributary to Anti-Aircraft Creek that is not consistent with any other mapped data sources nor our field observations. Tibbetts Creek flows north, crossing the project near its eastern terminus, and eventually flowing into Lake Sammamish. Schneider Creek crosses the project corridor near the northwestern terminus and eventually flows into Lake Sammamish. According to WDNR FPARS, Schneider Creek, Tibbetts Creek and Anti-Aircraft Creek are all considered fish bearing streams (Type F), and the unnamed stream is non-fish bearing (Type N) (WDNR 2017). SalmonScape has documented fish use within Tibbetts Creek by winter steelhead, kokanee, coho, and sockeye (WDFW 2017b). Additional WDFW SalmonScape data maps Anti-Aircraft Creek as having modeled distribution of winter steelhead, coho, sockeye and fall Chinook (WDFW 2017b).

City of Issaquah GIS Data Viewer (City of Issaquah 2017a) depicts three additional streams crossing the project corridor that are not mapped by WDNR FPARS or WDFW data. These additional streams are identified as 0169G, 0169I and 0169H. According to the City, these streams are all "Class 2 Streams with Salmonids" (2017). The stream hydrography dataset obtained from the City is depicted on Figure 2.

We also evaluated mapped wildlife habitats, which may be regulated under the IMC, to the extent they occur within protective stream and wetland buffers (IMC 18.10.360 B) and may also need to be documented as part of the NEPA/SEPA review processes. There are no priority wildlife habitats mapped within the project corridor and most habitats are outside of the Area of Investigation; the nearest priority wildlife habitats occur within Cougar Mountain Regional Wildland Park, which is located southwest of Newport Way. WDFW PHS maps this entire park area as a "Biodiversity Areas and Corridor" (WDFW 2017a). Within this park, WDFW PHS maps communal roosting habitat for Yuma myotis (*Myotis yumanensis*), Townsend's big-eared bat (*Corynorhinus townsendii*), and little brown bat (*Myotis lucifugus*)



(WDFW 2017a). There is an area of communal roosting habitat for these same bat species over 100 feet north of the project corridor and outside the Area of Investigation.

2.1.2. Soil Survey

The USDA-NRCS Web Soil Survey indicates six soil types within the project area (USDA-NRCS 2017a). Appendix A shows mapped soils within the project area. These mapped soil types are:

- Alderwood and Kitsap soils, very steep not hydric but may contain hydric inclusions
- Everett very gravelly sandy loam, 8 to 15 percent slopes not hydric and does not contain hydric inclusions
- Everett very gravelly sandy loam, 15 to 30 percent slopes not hydric but may contain hydric inclusions
- Kitsap silt loam, 2 to 8 percent slopes hydric
- Mixed alluvial land not hydric and does not contain hydric inclusions
- Sammamish silt loam hydric

Kitsap silt loam, 2 to 8 percent slopes and Sammamish silt loam are listed on the national hydric soils list (USDA-NRCS 2017b). Alderwood and Kitsap soils, very steep, Everett very gravelly sandy loam, 15 to 30 percent slopes are not hydric but may contain hydric inclusions (USDA-NRCS 2017b). Everett very gravelly sandy loam, 8 to 15 percent slopes, and mixed alluvial are not hydric and contain no hydric inclusions (USDA-NRCS 2017b).

2.2. Literature Review

2.2.1. Gateway Apartments - Previous Critical Areas Study and Final Mitigation Plan

The Gateway site is adjacent to Newport Way and at the time of the survey was undergoing extensive construction activities. We were granted ROE at the site and evaluated the remaining habitat within the area of investigation. Prior to construction, a Critical Areas Study and Final Mitigation Plan was developed for the Gateway project (Talasaea 2016). All proposed mitigation actions are outside of the Newport Way project area but are related to enhancement of Schneider Creek, its associated buffer and adjacent wetlands, as well as Tibbett's Creek and associated wetlands and buffers.

2.2.2. Riva Townhomes Site Plans

The Riva site includes one large parcel located on the east side of Newport way, across from the intersection with NW Oakcrest Drive (Figure 2). A review of the site plans prepared by Core Design (2015) indicates two stream features within this parcel. Anti-Aircraft Creek, which is located towards the northwest limits of the Riva property, has been recently relocated from its original channel into a new, more direct crossing of Newport Way (PBS 2017). The road crossing of Anti-Aircraft Creek was completed by the City as a separate project as part of the Riva Townhomes development agreement. The original stream channel was abandoned and converted to a storm drainage easement, and a new channel constructed at the replacement road crossing. This project has been completed separately from the Newport Way project but was designed to accommodate the new road section. A second stream, unnamed (identified as 0169H), is located on the southern portion of the Riva property, has been identified as Type F, and requires a culvert extension to accommodate frontage improvements, which is currently in construction by the Riva developers.



2.2.3. Former Bergsma Subdivision - Previous Culvert Survey

The former Bergsma site, now owned by City of Issaquah, is located on the southwest side of Newport Way in the southeastern portion of the project corridor near the intersection with SR 900 (Figure 2). The upstream portion of two mapped streams (City of Issaquah 2017a), 0169G and 0169I, are within these parcels. The Bergsma owners were previously in the process of developing plans for a subdivision development at these parcels. As part of this work, they had developed preliminary stream ordinary high water mark (OHWM) delineation and site survey associated with this proposed development and had initiated dialog with regulatory agencies.

Based on an email exchange from June 2016, WDFW visited these parcels and supported the designation of 0169G as fish-bearing but indicated that further slope information was needed to determine the accessibility and designation of 0169I (Peace 2016). In August 2016, Confluence Environmental Company (CEC) characterized the slope of 0169I as greater than 16 percent (CEC 2016).

2.3. Field Methods

GeoEngineers' biologists completed critical area site investigations on: October 10 and 30, 2017; May 3, 2018; and July 2, 2019. GeoEngineers field investigation was limited to the project footprint as defined by KPG drawings as well as extending up to 100 feet into adjacent areas where ROEs have been obtained. One property (Riva) was unable to be assessed directly because ROE was denied (Figure 2). In this case, over-the-fence observations were made from the adjacent road ROW onto the adjacent private parcels and existing data provided in developer drawings were referenced in lieu of direct OHWM delineation.

2.3.1. Wetland Assessment

Delineation of aquatic critical areas (wetlands and streams) was conducted in accordance with guidelines presented in IMC Chapter 18.10 (Environmental Protection) and using the *United States Army Corps of Engineers (USACE) Wetlands Delineation Manual* (Environmental Laboratory 1987) and the *Regional Supplement to the USACE Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region* (USACE 2010).

Potential wetland areas were evaluated based upon three parameters: (1) hydrophytic vegetation (USACE 2016); (2) hydric soils (USDA-NRCS 2017b); and (3) wetland hydrology (Ecology 1997). The presence of all three parameters may result in a jurisdictional wetland. Wetlands are rated using the Washington State Wetland Rating System for Western Washington (Hruby 2014).

2.3.2. Stream Assessment

The OHWM is used to define the jurisdictional boundary of streams and rivers according to the Clean Water Act, Washington State Growth Management Act, Washington State Shoreline Management Act and local critical areas ordinances. Methods used to identify the OHWM were consistent with local, state and federal protocols, including Regulatory Guidance Letter No. 05-05 (USACE 2005) and the Washington State manual Determining the Ordinary High Water Mark on Streams in Washington State (Olson and Stockdale 2016). We also referenced the Washington Administrative Code (WAC) for the definition of OHWM (WAC 173-22-030 § 11). Stream rating and identification of the buffer width was conducted in accordance with guidelines presented in IMC 18.10.785.



The OHWM of streams were evaluated using direct field observation by examining breaks in the topography, drift lines, shifts in vegetation and signs of water marks, according to USACE protocol as referenced from Regulatory Guidance Letter (No. 05-05), Ordinary High Water Mark Identification, December 7, 2005. Where ROE was not provided, as well as for the recently constructed stream channel at Anti-Aircraft Creek, which did not exhibit OHWM indicators, GeoEngineers referenced the developer drawings, which included OHWM delineations conducted by others, and confirmed general accuracy of these delineations as observed from the ROW in lieu of field delineation. On the former Bergsma site, now owned by the City of Issaquah, GeoEngineers also obtained the OHWM delineations completed by the developer and field-verified the accuracy of these data.

Streams within the City are rated following the rating system outlined in IMC Chapter 18.10.780. These ratings differ from the WDNR water typing classifications. Table 1 compares these two stream rating systems. This report will reference both methods.

TABLE 1. STREAM RATING SYSTEMS

City of Issaquah Stream Rating System	DNR Water Typing Classification
Class 1	Type S
Class 2 With Salmonids	Type F
Class 2 Streams	Type F
Class 3 Streams	Type Ns
Class 4 Streams	Type Ns

2.3.3. Fish and Wildlife Habitat

Fish and wildlife species presence and habitat use of the project corridor was evaluated through a review of available literature as well as general field observations. Sources of literature consulted prior to the field investigation included possible wildlife habitat relationships as documented by the USFWS endangered and threatened species list for the project location (USFWS 2017b) and the WDFW PHS map data (WDFW 2017a). We focused primarily on the PHS data to identify if regulated wildlife species or habitat conservation areas exist on or adjacent to the property. The focus of this evaluation was to document potential wildlife habitat and to make direct observations of physical habitat features (snags, nests, burrows, trails, dens, etc.). Fish and wildlife habitat areas were assessed according to IMC.

3.0 FINDINGS

We identified and mapped wetland habitat, stream habitat, fish and wildlife habitat, and stormwater features within the area of investigation during our field assessment. Additional data was obtained from developer drawings, particularly where ROE was not provided (Figure 2). A total of one offsite wetland (Wetland A) and six jurisdictional streams (Tibbetts Creek, 0169G, 0169H, 0169I, Anti-Aircraft Creek and Schneider Creek) were identified within the project corridor. Photographs of these features are included as Appendix B, Site Photographs. Critical areas are shown on Figure 3 through 5, Critical Area Results.

A photographic record was collected during the field visit to document existing site conditions. Representative photos have been included in Appendix B.



3.1. Wetlands

We delineated the jurisdictional wetland boundary for one wetland (Wetland A) adjacent to the project corridor. This wetland is associated with Tibbetts Creek and is outside the project corridor but parallel to the road through the southeastern portion of the project and partially within the Area of Investigation. Six formal data sample plots were established to make wetland identifications (Appendix C, Sample Plot Data Form). The wetland boundary was subsequently surveyed by KPG as depicted on Figures 4 and 5. We rated the wetland using the Washington State Wetland Rating System for Western Washington (Hruby 2014) as specified in IMC Chapter 18.10.029 (Wetland identification and functional rating). Appendix D, Wetland Rating Form includes the wetland rating form. A complete summary of wetland information and description of Wetland A is included in Appendix E, Wetland and Stream Descriptions.

Wetland A was identified as a Category II wetland based on the current rating system (Hruby 2014). Based on current regulatory code (IMC 18.10.640), a Category II wetland receives a 100-foot standard buffer. The standard buffer width can be decreased to 75 feet, when conditions are met as described in IMC 18.10.650.

3.2. Stream Delineation

GeoEngineers identified six streams within the project area: Tibbetts Creek, 0169G, 0169H, 0169I, Anti-Aircraft Creek and Schneider Creek. The streams 0169G, 0169H and 0169I are not mapped by WDNR, but are identified by the City (City of Issaquah 2017). Tibbetts Creek, the downstream sections of 0169G and 0169I, the upstream section of 0169H, and Schneider Creek were delineated during the field investigation and the resulting OHWM limits are shown on Figures 3 through 5. OHWM delineations of the upstream sections of 0169G and 0169I that were completed by others and provided in developer drawings were field-verified by GeoEngineers during a detailed stream reconnaissance as depicted on Figures 3 through 5. The downstream section of 0169H was also obtained from developer drawings for the Riva parcel (PACE 2016) and was field-verified from the ROW. Upstream and downstream sections of Anti-Aircraft Creek, which has been recently reconstructed, were obtained from the design plans for the replacement crossing since reliable field indicators have not developed since construction.

The City maps Tibbetts Creek, 0169G, 0169H, 0169I, Anti-Aircraft Creek and Schneider Creek as Class 2 Streams with Salmonids. However, based on our field assessment (GeoEngineers 2019), Stream 0169I contains intermittent, seasonal flow and does not contain salmonid habitat, indicating it would be more appropriately classified as Class 3 Streams according to IMC 18.10.780. This classification was confirmed by WDFW (Miles Penk, pers. comm. 2019). IMC 18.10.785(C) indicates a required standard vegetation buffer of 100 feet for Class 2 streams and 50 feet for Class 3 streams. Recommended buffer widths for each stream are shown on Figure 3. A summary of relevant stream information and general description of each stream are provided in Appendix E.

3.3. Stormwater Features

During the field investigation several stormwater features, that included stormwater pond, ditches, swales and culverts were identified. These features did not meet the criteria for wetland or stream habitat and were not delineated. The roadside ditches throughout the project corridor were in good condition. The old stream bed of Anti-Aircraft Creek (shown on Figure 4) has been converted to a storm drain easement for the adjacent roadway.



3.4. Fish and Wildlife Habitat

We did not observe any wildlife habitat, including no existing nest sites of any sensitive bird species, within the project corridor during our field work.

4.0 SUMMARY

This Critical Areas Report has been prepared for KPG on the behalf of the City of Issaquah for the proposed Newport Way Improvement Project. The objective of this report is to provide critical area baseline information for use during design and permitting for the project.

One offsite wetland (Wetland A) and six streams (Tibbetts, 0169G, 0169H, 0169I, Anti-Aircraft and Schneider) were identified during the field investigation. Wetland A is a Category II wetland and requires a 100-foot buffer due to a habitat score of 9. Tibbetts, 0169G, 0169H, Anti-Aircraft and Schneider Creek are all mapped by the City as Class 2 Stream with Salmonids (Type F) that will require a 100-foot buffer. Stream 0169I has been reviewed for potential fish utilization and concurrence has been obtained from WDFW that it is non-fish-bearing. Therefore, Stream 0169I is classified as Class 3 (Type Ns) and requires a 50-foot buffer. The buffers for the wetland and streams are identified below in Table 2. After project designs are finalized, potential wetland and buffer impacts should be assessed and, if needed, avoidance, minimization and mitigation options should be evaluated. If potential wetland and/or stream impacts are identified, a Mitigation Plan and other development permits may be required.

TABLE 2. CRITICAL AREAS SUMMARY

Wetland/ Stream ID	Category/Type	Buffer (feet)
Wetland A	Category II	100 ¹
Tibbetts Creek	Type F/Class 2 with Salmonids	100 ²
0169G	Type F/Class 2 with Salmonids	1002
0169H	Type F/Class 2 with Salmonids	100 ²
01691	Type Ns/Class 3	50 ²
Anti-Aircraft Creek	Type F/Class 2 with Salmonids	100 ²
Schneider Creek	Type F/Class 2 with Salmonids	100 ²

Notes:

- 1. Standard buffer based on IMC 18.10.640
- 2. Standard buffer based on IMC 18.10.785

5.0 LIMITATIONS

GeoEngineers has prepared this Critical Areas Report in general accordance with the scope and limitations of our proposal. Within the limitations of scope, schedule and budget, our services have been executed in accordance with the generally accepted practices for wetland and stream delineation in this area at the time this report was prepared. No warranty or other conditions, express or implied, should be understood.

This report has been prepared for the exclusive use of the KPG, Inc., authorized agents and regulatory agencies following the described methods and information available at the time of the work. No other party



may rely on the product of our services unless we agree in advance to such reliance in writing. The information contained herein should not be applied for any purpose or project except the one originally contemplated.

The applicant is advised to contact all appropriate regulatory agencies (local, state and federal) prior to design or construction of any development to obtain necessary permits and approvals.

6.0 REFERENCES

- City of Issaquah. 2017a. Interactive GIS Data Viewer. Available at: http://products.issaquahwa.gov/html5viewerpublic/index.html?viewer=issypublic
- City of Issaquah. 2017b. Newport Way Culvert Inventory. Prepared March 16, 2017. Provided to GeoEngineers by City of Issaquah.
- Confluence Environmental Company (CEC). 2016. Bergsma Property Stream Typing. Prepared for GFK Consulting. August 29, 2016.
- Core Design. 2015. Existing Conditions Map, Riva Townhomes. Prepared October 16, 2015.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of Wetland and Deep Water Habitats of the United States. Performed for Office of Biological Services, Fish and Wildlife Service, U.S. Department of the Interior, Washington, D.C.
- Environmental Laboratory. 1987. "U.S. Army Corps of Engineers Wetlands Delineation Manual," Technical Report Y-87-1, U.S. Army Engineer Waterways Experiment Station, Vicksburg, Massachusetts.
- GeoEngineers, Inc. 2019. "Fish Habitat Assessment for Streams 0169G and 0169I, Newport Way Road Improvements, Issaquah, Washington." Memorandum to Sessyle Asato, PE, Kelsey Anderson, PE, and Kirk Smith, PE (KPG, Inc.) prepared by David Conlin, PWS, Adam Wright, PWS, and Joe Callaghan, PWS (GeoEngineers, Inc.). GEI File No. 0252-039-01. September 5, 2019.
- Hruby, T. Revised. 2014. Washington State Wetland Rating System for Western Washington 2014 Update. Publication #14-06-029. Olympia, WA; Washington Department of Ecology
- Issaquah Municipal Code (IMC). Chapter 18 Environmental Protection Critical Areas Ordinance.
- Olson, Dr. Patricia and Stockdale, Eric. 2016. Determining the Ordinary High Water Mark on Streams in Washington State (Second Review Draft). April 2008, revised October 2016.
- PACE Engineering (PACE). 2016. Existing Stream Plan and Profile. August 10, 2016.
- Peace, Angie. 2016. RE: Bergsma Newport Way Stream Culvert 0169G. Fish and Wildlife Habitat Biologist, Washington State Department of Fish and Wildlife.
- Penk, Miles. 2019. RE: Notes/Follow-up From WDFW Site Visit at Newport Way. Email Communication with Habitat Biologist with Washington Department of Fish and Wildlife and David Conlin, Senior Biologist with GeoEngineers. October 21, 2019.



- PBS Engineering and Environmental (PBS). 2017. City of Issaquah Anti-Aircraft Creek Culvert Replacement Project, Bid Set.
- Talasaea Consultants, Inc. 2016. Critical Areas Study and Final Mitigation Plan, Gateway Apartments, City Permit #: SDP15-00002, Issaquah, Washington. Prepared for: the Wolff Company.
- United States Army Corps of Engineers (USACE). 2005. Regulatory Guidance Letter No. 05-05 Ordinary High Water Mark Identification. Riley Don T. December 7, 2005.
- United States Army Corps of Engineers (USACE). 2010. Regional Supplement to the Corps of Engineers
 Wetland Delineation Manual: Western Mountains, Valleys, and Coast Region, ed. J.S. Wakeley, R.
 W. Lichvar, and C.V. Noble. ERDC/EL TR-10-3. Vicksburg, MS: U.S. Army Engineer Research and
 Development Center.
- United States Army Corp of Engineers (USACE). 2016. Western Mountains, Valleys, and Coast 2016 Regional Wetland Plant List, US Army Corp of Engineers, Cold Regions Research and Engineering Laboratory.


 Available at:
 https://www.codot.gov/programs/environmental/wetlands/nwpl_wmvc_2016v1.pdf
- United States Department of Agriculture National Resource Conservation Service (USDA-NRCS). 2017a. Web Soil Survey. http://websoilsurvey.nrcs.usda.gov/app/.
- United States Department of Agriculture National Resource Conservation Service (USDA-NRCS). 2017b.

 National Hydric Soils List by State. Available at: http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/use/hydric/
- United States Fish and Wildlife Service (USFWS). 2017a. Wetlands Mapper. http://www.fws.gov/wetlands/Data/mapper.html
- United States Fish and Wildlife Service (USFWS). 2017b. Official Species List. Consultation Tracking Number 01EWFW00-2018-SLI-0336.
- Washington State Administrative Code (WAC). 2007. WAC 173-22-030. Definitions. http://apps.leg.wa.gov/WAC/default.aspx?cite=173-22-030
- Washington Department of Fish and Wildlife (WDFW). 2017a. Priority Habitats and Species (PHS) Interactive Mapping. https://wdfw.wa.gov/mapping/phs/
- Washington Department of Fish and Wildlife (WDFW). 2017b. SalmonScape Application. http://wdfw.wa.gov/mapping/salmonscape/
- Washington State Department of Ecology (Ecology). 1997. Washington State Wetlands Identification and Delineation Manual. Ecology Publication #96-94. March 1997.
- Washington State Department of Natural Resources (WDNR). 2017. Forest Practices Application Review System Mapping Application. Available at: http://fortress.wa.gov/dnr/app1/fpars/viewer.htm












- assist in showing features discussed in an attached document. GeoEngineers, Inc. cannot guarantee the accuracy and content of electronic files. The master file is stored by GeoEngineers, Inc. and will serve as the official record of this communication.

Data Source: King County GIS, ESRI, 2020 Survey

Projection: NAD 1983 StatePlane Washington North FIPS 4601 Feet






Feet

Newport Way Improvements Project Issaquah, Washington



Figure 3





assist in showing features discussed in an attached document. GeoEngineers, Inc. cannot guarantee the accuracy and content of electronic files. The master file is stored by GeoEngineers, Inc. and will serve as the official record of this communication.

Data Source: King County GIS, ESRI, 2020 Survey

Projection: NAD 1983 StatePlane Washington North FIPS 4601 Feet

Wetland Sample Plot Locations Survey Wetland Edge

Area of Investigation

Wetland Buffer 100 ft

No Right of Entry

Approximate Wetland Edge


Stream Bed - As Built


City of Issaquah Mapped Stream

Stream Based on Survey











Feet

Newport Way Improvements Project Issaquah, Washington




Figure 5



APPENDIX AData Review Sources

Newport Way Improvements Project

City of Issaquah, WA GIS



Newport Way Improvements Project



December 18, 2017 Data current as of May 4, 2020

Wetlands

Estuarine and Marine Deepwater

Estuarine and Marine Wetland

Freshwater Emergent Wetland

Freshwater Forested/Shrub Wetland

Freshwater Pond

Lake

Other

Riverine

This map is for general reference only. The US Fish and Wildlife Service is not responsible for the accuracy or currentness of the base data shown on this map. All wetlands related data should be used in accordance with the layer metadata found on the Wetlands Mapper web site.



SOURCE DATASET: PHSPlusPublic Query ID: P200504152241

REPORT DATE: 05/04/2020 3.23

Common Name Scientific Name Notes	Site Name Source Dataset Source Record Source Date	Priority Area Occurrence Type More Information (URL) Mgmt Recommendations	Accuracy	Federal Status State Status PHS Listing Status	Sensitive Data Resolution	Source Entity Geometry Type	
Big brown bat		Communal Roost	GPS	N/A	Υ	WA Dept. of Fish and Wildlife	
Eptesicus fuscus	WS_OccurPoint 145195	Biotic detection		N/A	TOWNSHIP	Points	
	March 07, 2011	http://wdfw.wa.gov/publications/pub.php?		PHS LISTED			
Biodiversity Areas And	KING COUNTY (EASTSIDE)	Terrestrial Habitat	1/4 mile (Quarter	N/A	N	WA Dept. of Fish and Wildlife	
	PHSREGION 902805	N/A	`	N/A	AS MAPPED	Polygons	
		http://wdfw.wa.gov/publication	ns/pub.php?	PHS LISTED			
Coho	Tibbetts Creek	Breeding Area	NA	N/A	N		
Oncorhynchus kisutch	SWIFD	Breeding area		N/A	AS MAPPED	Lines	
	28259	http://wdfw.wa.gov/wlm/diversty/soc/soc.htm		PHS LISTED			
Coho	Tibbetts Creek	Occurrence	NA	Candidate	N	WDFW Fish Program	
Oncorhynchus kisutch	SASI	Occurrence		N/A	AS MAPPED	Lines	
	3120	http://wdfw.wa.gov/wlm/diversty/soc/soc.htm		PHS Listed			
Freshwater Emergent	N/A	Aquatic Habitat	NA	N/A	N	US Fish and Wildlife Service	
Ŭ	NWIWetlands	Aquatic habitat		N/A	AS MAPPED	Polygons	
		http://www.ecy.wa.		PHS Listed			
Freshwater Forested/Shrub	N/A	Aquatic Habitat	NA	N/A	N	US Fish and Wildlife Service	
	NWIWetlands	Aquatic habitat		N/A	AS MAPPED	Polygons	
		http://www.ecy.wa.		PHS Listed			
Freshwater Forested/Shrub	N/A	Aquatic Habitat	NA	N/A	N	US Fish and Wildlife Service	
	NWIWetlands	Aquatic habitat		N/A	AS MAPPED	Polygons	
				PHS Listed			

05/04/2020 3.23

Common Name Scientific Name Notes	Site Name Source Dataset Source Record Source Date	Priority Area Occurrence Type More Information (URL) Mgmt Recommendations	Accuracy	Federal Status State Status PHS Listing Status	Sensitive Data Resolution	Source Entity Geometry Type
Kokanee Oncorhynchus nerka	Tibbetts Creek SWIFD 28262	Occurrence/Migration N Occurrence/migration http://wdfw.wa.gov/wlm/diversty/	IA /soc/soc.htm	N/A N/A PHS LISTED	N AS MAPPED	Lines
Little Brown Bat Myotis lucifugus	WS_OccurPoint 145125 July 10, 2017	Breeding Area G Biotic detection http://wdfw.wa.gov/publications/	GPS	N/A N/A PHS LISTED	Y TOWNSHIP	WA Dept. of Fish and Wildlife Points
Little Brown Bat Myotis lucifugus	WS_OccurPoint 148355 June 20, 2018		GPS	N/A N/A PHS LISTED	Y TOWNSHIP	WA Dept. of Fish and Wildlife Points
Resident Coastal Cutthroat Oncorhynchus clarki	Tibbetts Creek SWIFD 28256	Occurrence/Migration N Occurrence/migration http://wdfw.wa.gov/wlm/diversty/	JA /soc/soc.htm	N/A N/A PHS LISTED	N AS MAPPED	Lines
Sockeye Oncorhynchus nerka	Tibbetts Creek SWIFD 28263	Occurrence/Migration N Occurrence/migration http://wdfw.wa.gov/wlm/diversty/	IA /soc/soc.htm	N/A N/A PHS LISTED	N AS MAPPED	Lines
Sockeye Oncorhynchus nerka	Tibbetts Creek SASI 5200	Occurrence N Occurrence http://wdfw.wa.gov/wlm/diversty/	JA /soc/soc.htm	Not Warranted N/A PHS Listed	N AS MAPPED	WDFW Fish Program Lines
Steelhead Oncorhynchus mykiss	Tibbetts Creek SASI 6154	Occurrence N Occurrence http://wdfw.wa.gov/wlm/diversty/	IA /soc/soc.htm	Threatened N/A PHS Listed	N AS MAPPED	WDFW Fish Program Lines
Townsend's Big-eared Bat Corynorhinus townsendii	WS_OccurPoint 109377 March 07, 2011	Communal Roost G Biotic detection http://wdfw.wa.gov/publications/	GPS /pub.php?	N/A Candidate PHS LISTED	Y TOWNSHIP	WA Dept. of Fish and Wildlife Points


05/04/2020 3.23


Common Name Scientific Name	Site Name Source Dataset Source Record	Priority Area Occurrence Type More Information (URL)	Accuracy	Federal Status State Status	Sensitive Data Resolution	Source Entity Geometry Type
Notes	Source Date	Mgmt Recommendations		PHS Listing Status		
140103	200.00 20.0					
Townsend's Big-eared Bat		Communal Roost	GPS	N/A	Υ	WA Dept. of Fish and Wildlife
Corynorhinus townsendii	WS_OccurPoint 112189	Biotic detection		Candidate	TOWNSHIP	Points
	February 26, 2012	http://wdfw.wa.gov/publicati	ons/pub.php?	PHS LISTED		
Townsend's Big-eared Bat		Communal Roost	GPS	N/A	Υ	WA Dept. of Fish and Wildlife
Corynorhinus townsendii	WS_OccurPoint 145143	Biotic detection		Candidate	TOWNSHIP	Points
	May 04, 2016	http://wdfw.wa.gov/publications/pub.php?		PHS LISTED		
Wetlands	ISSAQUAH CREEK	Aquatic Habitat	1/4 mile (Quarter	N/A	N	WA Dept. of Fish and Wildlife
	PHSREGION 902512	N/A	`	N/A	AS MAPPED	Polygons
		http://www.ecy.wa.		PHS LISTED		
Winter Steelhead	Tibbetts Creek	Breeding Area	NA	N/A	N	
Oncorhynchus mykiss	SWIFD	Breeding area		N/A	AS MAPPED	Lines
	28266	http://wdfw.wa.gov/wlm/diversty/soc/soc.htm		PHS LISTED		
Yuma myotis		Breeding Area	GPS	N/A	Υ	WA Dept. of Fish and Wildlife
Myotis yumanensis	WS_OccurPoint 145124	Biotic detection	0.0	N/A	TOWNSHIP	Points
	July 10, 2017	http://wdfw.wa.gov/publicati	ons/pub.php?	PHS LISTED		
Yuma myotis		Breeding Area	GPS	N/A	Υ	WA Dept. of Fish and Wildlife
Myotis yumanensis	WS_OccurPoint 148356	Biotic detection		N/A	TOWNSHIP	Points
	June 20, 2018	http://wdfw.wa.gov/publicati	ons/pub.php?	PHS LISTED		


DISCLAIMER. This report includes information that the Washington Department of Fish and Wildlife (WDFW) maintains in a central computer database. It is not an attempt to provide you with an official agency response as to the impacts of your project on fish and wildlife. This information only documents the location of fish and wildlife resources to the best of our knowledge. It is not a complete inventory and it is important to note that fish and wildlife resources may occur in areas not currently known to WDFW biologists, or in areas for which comprehensive surveys have not been conducted. Site specific surveys are frequently necessary to rule out the presence of priority resources. Locations of fish and wildlife resources are subject to vraition caused by disturbance, changes in season and weather, and other factors. WDFW does not recommend using reports more than six months old.

05/04/2020 3.23

WDFW Test Map







Source: Esri, DigitalGlobe, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community



NRCS

Natural Resources Conservation Service A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for King County Area, Washington

Newport Way Improvements Project



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2 053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	
Soil Map	
Soil Map	9
Legend	
Map Unit Legend	
Map Unit Descriptions	
King County Area, Washington	13
AkF—Alderwood and Kitsap soils, very steep	13
Bh—Bellingham silt loam	14
EvC—Everett very gravelly sandy loam, 8 to 15 percent slopes	15
EvD—Everett very gravelly sandy loam, 15 to 30 percent slopes	16
KpB—Kitsap silt loam, 2 to 8 percent slopes	18
Ma—Mixed alluvial land	19
Sh—Sammamish silt loam	
Sk—Seattle muck	21
References	

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



MAP LEGEND

Area of Interest (AOI)

Area of Interest (AOI)

Soils

Soil Map Unit Polygons

Soil Map Unit Lines

Soil Map Unit Points

Special Point Features

(o)

Blowout

Borrow Pit

Clay Spot

Closed Depression

Gravel Pit

Gravelly Spot

Landfill Lava Flow

Marsh or swamp

Mine or Quarry

Miscellaneous Water Perennial Water

Rock Outcrop

Saline Spot

Sandy Spot

Severely Eroded Spot

Sinkhole

Slide or Slip

Sodic Spot

å

Spoil Area

Stony Spot

Very Stony Spot

Ŷ

Wet Spot Other

Δ

Special Line Features

Water Features

Streams and Canals

Transportation

Rails

Interstate Highways

US Routes

Major Roads

00

Local Roads

Background

Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:24.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: King County Area, Washington Survey Area Data: Version 15, Sep 16, 2019

Soil map units are labeled (as space allows) for map scales 1:50.000 or larger.

Date(s) aerial images were photographed: Jul 1, 2019—Jul 25, 2019

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
AkF	Alderwood and Kitsap soils, very steep	16.7	25.9%
Bh	Bellingham silt loam	0.5	0.7%
EvC	Everett very gravelly sandy loam, 8 to 15 percent slopes	19.5	30.3%
EvD	Everett very gravelly sandy loam, 15 to 30 percent slopes	6.1	9.5%
КрВ	Kitsap silt loam, 2 to 8 percent slopes	8.3	12.9%
Ма	Mixed alluvial land	6.5	10.1%
Sh	Sammamish silt loam	5.7	8.9%
Sk	Seattle muck	1.1	1.8%
Totals for Area of Interest	,	64.5	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not

mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

King County Area, Washington

AkF—Alderwood and Kitsap soils, very steep

Map Unit Setting

National map unit symbol: 1hmsn

Elevation: 50 to 800 feet

Mean annual precipitation: 25 to 60 inches Mean annual air temperature: 48 to 52 degrees F

Frost-free period: 160 to 220 days

Farmland classification: Not prime farmland

Map Unit Composition

Alderwood and similar soils: 50 percent Kitsap and similar soils: 25 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Alderwood

Setting

Landform: Moraines, till plains

Parent material: Basal till with some volcanic ash

Typical profile

H1 - 0 to 12 inches: gravelly ashy sandy loam H2 - 12 to 27 inches: very gravelly sandy loam H3 - 27 to 60 inches: very gravelly sandy loam

Properties and qualities

Slope: 25 to 70 percent

Depth to restrictive feature: 24 to 40 inches to densic material

Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.06 in/hr)

Depth to water table: About 18 to 37 inches

Frequency of flooding: None Frequency of ponding: None

Available water storage in profile: Very low (about 2.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: B Hydric soil rating: No

Description of Kitsap

Setting

Landform: Terraces

Parent material: Lacustrine deposits with a minor amount of volcanic ash

Typical profile

H1 - 0 to 5 inches: ashy silt loam H2 - 5 to 24 inches: ashy silt loam

H3 - 24 to 60 inches: stratified silt to silty clay loam

Properties and qualities

Slope: 25 to 70 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: None Frequency of ponding: None

Available water storage in profile: High (about 11.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7e

Hydrologic Soil Group: C Hydric soil rating: No

Bh—Bellingham silt loam

Map Unit Setting

National map unit symbol: 1hmsw

Elevation: 0 to 820 feet

Mean annual precipitation: 35 to 60 inches Mean annual air temperature: 50 degrees F

Frost-free period: 150 to 210 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Bellingham and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Bellingham

Setting

Landform: Depressions, drainageways

Parent material: Alluvium

Typical profile

H1 - 0 to 11 inches: silt loam
H2 - 11 to 60 inches: silty clay loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 0 to 12 inches

Frequency of flooding: None

Frequency of ponding: None

Available water storage in profile: Very high (about 12.6 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5w

Hydrologic Soil Group: C/D

Forage suitability group: Wet Soils (G002XN102WA)

Hydric soil rating: Yes

Minor Components

Alderwood

Percent of map unit: 5 percent

Hydric soil rating: No

Everett

Percent of map unit: 5 percent

Hydric soil rating: No

Seattle

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

EvC—Everett very gravelly sandy loam, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2t62b

Elevation: 30 to 900 feet

Mean annual precipitation: 35 to 91 inches
Mean annual air temperature: 48 to 52 degrees F

Frost-free period: 180 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Everett and similar soils: 80 percent Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Everett

Setting

Landform: Kames, eskers, moraines

Landform position (two-dimensional): Shoulder, footslope Landform position (three-dimensional): Crest, base slope

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Sandy and gravelly glacial outwash

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

A - 1 to 3 inches: very gravelly sandy loam
Bw - 3 to 24 inches: very gravelly sandy loam
C1 - 24 to 35 inches: very gravelly loamy sand
C2 - 35 to 60 inches: extremely cobbly coarse sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Somewhat excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water storage in profile: Low (about 3.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4s

Hydrologic Soil Group: A

Forage suitability group: Droughty Soils (G002XN402WA), Droughty Soils

(G002XS401WA), Droughty Soils (G002XF403WA)

Hydric soil rating: No

Minor Components

Indianola

Percent of map unit: 10 percent Landform: Eskers, kames, terraces

Landform position (three-dimensional): Riser

Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: No

Alderwood

Percent of map unit: 10 percent

Landform: Ridges, hills

Landform position (two-dimensional): Shoulder

Landform position (three-dimensional): Nose slope, talf

Down-slope shape: Linear, convex Across-slope shape: Convex

Hydric soil rating: No

EvD—Everett very gravelly sandy loam, 15 to 30 percent slopes

Map Unit Setting

National map unit symbol: 2t62c

Elevation: 30 to 900 feet

Mean annual precipitation: 35 to 91 inches
Mean annual air temperature: 48 to 52 degrees F

Frost-free period: 180 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Everett and similar soils: 80 percent Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Everett

Setting

Landform: Kames, eskers, moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Convex

Parent material: Sandy and gravelly glacial outwash

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

A - 1 to 3 inches: very gravelly sandy loam
Bw - 3 to 24 inches: very gravelly sandy loam
C1 - 24 to 35 inches: very gravelly loamy sand
C2 - 35 to 60 inches: extremely cobbly coarse sand

Properties and qualities

Slope: 15 to 30 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Somewhat excessively drained

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water storage in profile: Low (about 3.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: A

Forage suitability group: Droughty Soils (G002XN402WA), Droughty Soils

(G002XS401WA) *Hydric soil rating:* No

Minor Components

Alderwood

Percent of map unit: 10 percent

Landform: Ridges, hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope, nose slope, talf

Down-slope shape: Linear, convex Across-slope shape: Convex

Hydric soil rating: No

Indianola

Percent of map unit: 10 percent Landform: Eskers, kames, terraces

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: No

KpB—Kitsap silt loam, 2 to 8 percent slopes

Map Unit Setting

National map unit symbol: 1hmt9

Elevation: 0 to 590 feet

Mean annual precipitation: 37 inches Mean annual air temperature: 50 degrees F

Frost-free period: 160 to 200 days

Farmland classification: All areas are prime farmland

Map Unit Composition

Kitsap and similar soils: 85 percent *Minor components:* 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Kitsap

Settina

Landform: Terraces

Parent material: Lacustrine deposits with a minor amount of volcanic ash

Typical profile

H1 - 0 to 5 inches: silt loam H2 - 5 to 24 inches: silt loam

H3 - 24 to 60 inches: stratified silt to silty clay loam

Properties and qualities

Slope: 2 to 8 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Moderately well drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.06 to 0.20 in/hr)

Depth to water table: About 18 to 36 inches

Frequency of flooding: None Frequency of ponding: None

Available water storage in profile: High (about 11.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3w

Hydrologic Soil Group: C

Forage suitability group: Soils with Few Limitations (G002XN502WA)

Hydric soil rating: No

Minor Components

Alderwood

Percent of map unit: 10 percent

Hydric soil rating: No

Bellingham

Percent of map unit: 3 percent Landform: Depressions Hydric soil rating: Yes

Seattle

Percent of map unit: 1 percent Landform: Depressions Hydric soil rating: Yes

Tukwila

Percent of map unit: 1 percent Landform: Depressions Hydric soil rating: Yes

Ma—Mixed alluvial land

Map Unit Setting

National map unit symbol: 1hmtf

Elevation: 0 to 590 feet

Mean annual precipitation: 25 to 90 inches Mean annual air temperature: 46 to 54 degrees F

Frost-free period: 160 to 200 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Alluvial land, mixed, and similar soils: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Alluvial Land, Mixed

Typical profile

H1 - 0 to 8 inches: sand H2 - 8 to 20 inches: fine sand H3 - 20 to 60 inches: sand

H4 - 60 to 70 inches: loamy fine sand, gravelly sand

H4 - 60 to 70 inches:

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Natural drainage class: Well drained

Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95

to 19.98 in/hr)

Depth to water table: About 12 to 36 inches

Frequency of flooding: Frequent Frequency of ponding: None

Available water storage in profile: Very low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: A Hydric soil rating: No

Sh—Sammamish silt loam

Map Unit Setting

National map unit symbol: 1hmv3

Elevation: 0 to 50 feet

Mean annual precipitation: 45 to 60 inches

Mean annual air temperature: 48 to 52 degrees F

Frost-free period: 180 to 220 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Sammamish and similar soils: 84 percent

Minor components: 16 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sammamish

Setting

Landform: Flood plains
Parent material: Alluvium

Typical profile

H1 - 0 to 12 inches: silt loam

H2 - 12 to 60 inches: stratified loamy sand to silt loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Somewhat poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to

high (0.57 to 1.98 in/hr)

Depth to water table: About 12 to 24 inches

Frequency of flooding: Occasional Frequency of ponding: None

Available water storage in profile: High (about 11.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: B/D

Forage suitability group: Seasonally Wet Soils (G002XN202WA)

Hydric soil rating: Yes

Minor Components

Bellingham

Percent of map unit: 10 percent

Landform: Depressions Hydric soil rating: Yes

Puget

Percent of map unit: 5 percent Landform: Depressions Hydric soil rating: Yes

Puyallup

Percent of map unit: 1 percent

Hydric soil rating: No

Sk—Seattle muck

Map Unit Setting

National map unit symbol: 1hmv4

Elevation: 0 to 1,000 feet

Mean annual precipitation: 25 to 50 inches Mean annual air temperature: 48 to 52 degrees F

Frost-free period: 150 to 250 days

Farmland classification: Prime farmland if drained

Map Unit Composition

Seattle and similar soils: 75 percent *Minor components:* 25 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Seattle

Setting

Landform: Depressions

Parent material: Grassy organic material

Typical profile

H1 - 0 to 11 inches: muck

H2 - 11 to 60 inches: stratified mucky peat to muck

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches Natural drainage class: Very poorly drained

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to

high (0.57 to 1.98 in/hr)

Depth to water table: About 0 inches

Frequency of flooding: None Frequency of ponding: Frequent

Available water storage in profile: Very high (about 23.5 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 5w

Hydrologic Soil Group: B/D

Forage suitability group: Wet Soils (G002XN102WA)

Hydric soil rating: Yes

Minor Components

Shalcar

Percent of map unit: 10 percent

Landform: Depressions Hydric soil rating: Yes

Tukwila

Percent of map unit: 10 percent

Landform: Depressions Hydric soil rating: Yes

Bellingham

Percent of map unit: 3 percent Landform: Depressions Hydric soil rating: Yes

Norma

Percent of map unit: 2 percent Landform: Depressions Hydric soil rating: Yes

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf

APPENDIX BSite Photographs



Photograph 1. Tibbetts Creek viewed from the southern side of Newport Way near intersection of Newport Way and State Route 900 (Photo taken October 30, 2017)



Photograph 2. Tibbetts Creek near intersection with Newport Way and State Route 900, viewed downstream from the north side of Newport Way (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 3. Box culvert under Newport Way for Tibbetts Creek (Photo taken October 30, 2017).



Photograph 4. Stream 0169G viewed to the northeast (downstream) from Newport Way from the culvert outlet (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 5. Stream 0169G viewed further downstream of Newport Way, illustrating modified channel, bank and adjacent residential property on the left (Photo taken October 30, 2016).



Photograph 6. Stream 0169G outlet on the left, viewed looking back towards Newport Way (Photo taken July 28, 2016).

Newport Way Improvements Project Issaquah, Washington





Photograph 7. Wetland A interior, including Tibbetts Creek (Photo taken May 3, 2018).



Photograph 8. Wetland A buffer conditions as viewed from Newport Way. Wetland A and Tibbetts creek are in background (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 9. Stream 0169I viewed at culvert outlet on downstream (northeast) side of Newport Way (Photo taken April 27, 2018).



Photograph 10. Stream 0169I viewed looking upstream from Newport Way (Photo taken July 2, 2019).

Newport Way Improvements Project Issaquah, Washington





Photograph 11. Typical vegetation conditions adjacent to Newport Way, viewed to the southeast from near Stream 0169I (Photo taken October 30, 2017).



Photograph 12. Stream 0169H upstream channel conditions (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 13. Stream 0169H at culvert outlet on north side of Newport Way (Photo taken October 30, 2017).



Photograph 14. Stream 0169H viewed to the north from Newport Way, depicting degraded buffer conditions in the foreground (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 15. Recently reconstructed channel of Anti-Aircraft Creek (Photo taken October 10, 2017).



Photograph 16. Prior channel of Anti-Aircraft Creek downstream of Newport Way. This channel has been replaced with a new alignment and newly constructed channel and has been converted to a storm drainage easement (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 17. Typical roadside conditions along Newport Way near Sammamish Point and Gateway development projects (Photo taken October 30, 2017).



Photograph 18. Typical roadside conditions along Newport Way near Sammamish Point and Gateway development projects (Photo taken October 30, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 19. Schneider Creek corridor upstream (south) of Newport Way (Photo taken October 10, 2017).



Photograph 20. Schneider Creek culvert inlet and riparian vegetation viewed from Newport Way (Photo taken October 10, 2017).

Newport Way Improvements Project Issaquah, Washington





Photograph 21. Schneider Creek Newport Way crossing culvert outlet viewed upstream toward Newport Way from the northeast. The buffer consists of mowed grass on this residential property. (Photo taken October 10, 2017)



Photograph 22. Schneider Creek viewed northeast (downstream) from offsite portion of stream corridor (Photo taken October 10, 2017).

Newport Way Improvements Project Issaquah, Washington



APPENDIX CSample Plot Data Form

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site: Newport Way Improvement Applicant/Owner: City of Issaquah		Improvements Project		City/County:	King		Sampling Date: 5/3/2018			5/3/2018
		ıh			State	: WA		Sampling	Point: SP1	L
Investigator(s):	DBC	_		Section/Townsh	ip/Range:	S29 T24N R06E				
Landform (hillslope, terr	race, etc.):	terrace		Local Relief (con	ıcave, convex, r	none):	concave	Slop	oe (%): <u><5%</u>	%
Subregion (LLR):	Α		_ Lat:	47.545523°	Long	: <u>-122.067415°</u>	Datum:	WGS 1984		
Soil Map Unit Name:	Alderwood an	d Kitsap soils, v	very steep		_ N	WI Classification:	NA			
Are climatic/hydrologic	conditions on the	e site typical fo	or this time of year?		✓ Yes	☐ No	(if no, explain	in Remarks.)		
Are Vegetation	Are Vegetation Soil Hydrology significantly			rbed?	Are "normal	circumstances" pre	present?			
Are	Soil H	Hydrology	naturally problem	natic?	(if needed, ex	xplain any answers	in Remarks.)			
SUMMARY OF FIND				1						ı
Hydrophytic Vegetation Hydric Soil Present? Weltand Hydrology Pres		✓ Yes ✓ Yes ✓ Yes	No No No	Is the sampled a Wetland?	irea within a	✓ Yes				
Remarks:										
VEGETATION - Use	scientific nam	nes of plants	5.							
Tree Stratum_		•	Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test	Worksheet:			
1. Alnus rubra			20	Y	FAC	Number of domi	nant Species			
2.						That are OBL, FA		4		(A)
3.			-			4				
4.			20	- Total Cavar		Total Number of		4		(D)
Sapling/Shurb Strat	um		20	= Total Cover		Species Acro	ss All Strata:	4		(B)
1. Rubus spectabilis	um_		15	٧	FAC	Percent of domir	ant Species			
2. Salix sp.			25	Y	FACW	That are OBL, FA		10	0	(A/B)
<i>3</i> .										
4.						Prevalence Index				
5.				T. I. I. C.	_	Total % Co		Multiply by:		
Iloub Ctuatuus			40	= Total Cover		OBL Species	0	x 1 =	0	
Herb Stratum 1. Athyrium felix-femine	~		5	N	FAC	FACW Species FAC Species	95	x 2 = x 3 =	190 150	
2. Equisetum arvense	<u> </u>		10	N	FAC	FACU Species		x 4 =	0	
3. Phalaris arundinacea	1		70	Y	FACW	UPL Species		x 5 =	0	
4.			_			Column Totals:	145	(A)	340	(B)
5.								5.44		
<i>6.</i> <i>7</i> .			_			Prev	alence Index :	= B/A =	2.34	
8.			_		-	Hydrophytic Veg	etation Indica	tors:		
9.						1 - Rapid Tes				
10.						2 - Dominano				
11.			_		_	✓ 3 - Prevalence				
			85	= Total Cover		4 - Morpholo	ogical Adaptat	ions¹ (provide su	pporting d	ata in
Woody Vine Stratur	<u>n</u> _					Remarks or o	•			
1.						5 - Wetland I	Non-Vascular	Plants ¹		
2.						Problem Hyd	Irophytic Vege	etation (Explain)		
			0	= Total Cover		¹ Indicators of	hydric soil	and wetland h	ydrology	y must
% Bare Ground	in Herb Stratum	ı	_			be present, ur	nless distur	bed or proble	matic.	
Remarks:						Hydrophytic V	egetation Pre	sent?	es 🔲 N	lo
						•				

SOIL Sampling Point: SP1 Depth Matrix **Redox Features** Type^1 Loc² (inches) Color (moist) Texture Remarks Color (moist) % 10 YR 3/2 0 to 6 6 to 20 10 YR 3/1 85 5 YR 4/6 15 С PL loam ¹Type: C=Concentration, D=Depletion, RM-Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix Indicators for Problematic Hydric Soils³: Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.) ☐ Histisol (A1) Sandy Redox (S5) 2 cm Muck (A10) Histic Epipedon (A2) Stripped Matrix (S6) Red Parent Material (TF2) Black Histic (A3) Loamy Mucky Mineral (F1) (except MLRA 1) ☐ Very Shallow Dard Surface (TF12) Loamy Gleyed Matrix (F2) Hydrogen Sulfide (A4) Other (Explain in Remarks) Depleted Below Dark Surface (A11) Depleted Matrix (F3) maicators or myarophytic vegetation Thick Dark Surface (A12) Redox Dark Surface (F6) and wetland hydrology must be present, Sandy Mucky Mineral (S1) Depleted Dark Surface (F7) unless disturbed or problematic. Sandy Gleyed Matrix (S4) Redox Depressions (F8) Restrictive Layer (if present): Hydric Soil Present? Type: Depth (inches): Remarks: **HYDROLOGY** Wetland Hydrology Indicators: Primary Indicators (minimum of one required; check all that apply) Secondary Indicators (2 or more required) Surface Water (A1) ■ Water-Stained Leaves (B9) (except MLRA Water-Stained Leaves (B9) (MLRA High Water Table (A2) 1, 2, 4A, and 4B) 1, 2, 4A, and 4B) Saturation (A3) Salt Crust (B11) Drainage Patterns (B10) Water Marks (B1) Aquatic Invertebrates (B13) Dry-Season Water Table (C2) Sediment Deposits (B2) Hydrogen Sulfide Odor (C1) Saturated Visible on Aerial Imagery (C9) Drift Deposits (B3) Oxidized Rhizospheres along Living Roots (C3) Geomorphic Position (D2) Algal Mat or Crust (B4) Presence of Reduction Iron (C4) Shallow Aquitard (D3) Iron Deposits (B5) Recent Iron Reduction Tilled Soils (C6) ✓ FAC-Neutral Test (D5) Stunted or Stressed Plants (D1) (LRR A) Raised Ant Mounds (D6) (LRR A) Surface Soil Cracks (B6) Other (Explain in Remarks) Frost-Heave Hummocks (D7) Inundation Visible on Aerial Imagery (B7) Sparsely Vegetated Concave Surface (B8) Field Observations: Wetland Hydrology Present? Surface Water Present? ✓ No Depth (inches): \neg_{Yes} Water Table Present? Depth (inches): ✓ Yes ☐ No Yes ✓ No Saturation Present? Yes No Depth (inches): (includes capillary fringe)

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Remarks: two secondary indicators

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site:	Newport Way Improvement	nts Project	_City/County:	King		Sampling Date: 5/3/2018 Sampling Point: SP2		
Applicant/Owner:	City of Issaquah	of Issaquah		State	: WA			
Investigator(s):	DBC		Section/Townsh	ip/Range:	S29 T24N R06E			
Landform (hillslope, terr	ace, etc.): terrace		Local Relief (con	cave, convex, r	none): <u>c</u>	oncave	Slope (%): <u><</u>	5%
Subregion (LLR):	Α	Lat:	47.545410°	Long	: <u>-122.067498°</u>	Datum: V	VGS 1984	
Soil Map Unit Name:	Alderwood and Kitsap soils	s, very steep		_ N	WI Classification: N	IA		
Are climatic/hydrologic	conditions on the site typical	for this time of year?	•	✓ Yes	□ No (i	f no, explain in	Remarks.)	
Are Vegetation	Soil Hydrology	significantly distu	irbed?	Are "normal	circumstances" pres	present? Yes No		
Are Vegetation	Soil Hydrology	naturally problen	natic?	(if needed, ex	xplain any answers i	n Remarks.)		
SUMMARY OF FIND			.					
Hydrophytic Vegetation Hydric Soil Present? Weltand Hydrology Pres	Yes	☐ No ✓ No ✓ No	Is the sampled a Wetland?	rea within a	Yes Vo			
Remarks: FAC veg comm	nunity meets hydrophytic veį	getation standard; no	other parameters	met.				
VEGETATION - Use s	scientific names of plar	nts.						
Tree Stratum		Absolute % Cover	Dominant Species?	Indicator	Dominance Test \	Worksheet:		
1. Alnus rubra		60	Species? Y	Status FAC	Number of domin	ant Species		
2.					That are OBL, FAC	CW, or FAC:	4	(A)
3.								
4.		60	= Total Cover		Total Number of I Species Acros		5	(B)
Sapling/Shurb Stratu	um		_		- Species 7 to 65			(5)
1. Rubus spectabilis	- <u></u>	30	Υ	FAC	Percent of domina	ant Species		
2. Symphoricarpos albus	S	30	Υ	FACU	That are OBL, FAC	CW, or FAC:	80	(A/B)
3.						NA (
<u>4.</u> 5.					Prevalence Index Total % Cov		/ultiply by:	
<u>. </u>		60	= Total Cover	-	OBL Species	0 x		
Herb Stratum			_		FACW Species	0 x	2 = 0	
1. Polystichum munitum	1	20	Υ	FAC	FAC Species	150 x	3 = 450	
2. Athyrium felix-femino	7	30	Υ	FAC	FACU Species	30 x		
3. Equisetum arvense		10	N	FAC	UPL Species	0 x 180 (/		(D)
4. 5.		_ -		-	Column Totals:	180 (7	A)	(B)
6. 7.					Preva	lence Index = B	3/A = 3.17	
]			
8.					Hydrophytic Vege			
9. 10.					1 - Rapid Test ✓ 2 - Dominance		c vegetation	
11.				-	3 - Prevalence		L	
		60	= Total Cover		4 - Morpholog	gical Adaptation	ns ¹ (provide supporting	data in
Woody Vine Stratun	n		_			a separate she		
	_				5 - Wetland N	•		
<u>1.</u> 2.			-		Problem Hydr			
		0	= Total Cover		-		id wetland hydrolo	gy must
% Bare Ground	in Herb Stratum	_	-			•	ed or problematic.	
Remarks: primarly FAC	community				Hydrophytic Ve	getation Preser	nt?	No
							•	

SOIL Sampling Point: SP2 Depth Matrix **Redox Features** Type¹ Loc² Texture Remarks (inches) Color (moist) Color (moist) 0-12" 10YR 2/2 100 loam ¹Type: C=Concentration, D=Depletion, RM-Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix Indicators for Problematic Hydric Soils³: Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.) Histisol (A1) Sandy Redox (S5) 2 cm Muck (A10) Histic Epipedon (A2) Stripped Matrix (S6) Red Parent Material (TF2) Black Histic (A3) Loamy Mucky Mineral (F1) (except MLRA 1) ☐ Very Shallow Dard Surface (TF12) Loamy Gleyed Matrix (F2) Hydrogen Sulfide (A4) Other (Explain in Remarks) Depleted Below Dark Surface (A11) Depleted Matrix (F3) maicators or myarophytic vegetation Thick Dark Surface (A12) Redox Dark Surface (F6) and wetland hydrology must be present, Sandy Mucky Mineral (S1) Depleted Dark Surface (F7) unless disturbed or problematic. Sandy Gleyed Matrix (S4) Redox Depressions (F8) Restrictive Layer (if present): Hydric Soil Present? Type: Yes Vo Depth (inches): Remarks: **HYDROLOGY** Wetland Hydrology Indicators: Primary Indicators (minimum of one required; check all that apply) Secondary Indicators (2 or more required) Surface Water (A1) ■ Water-Stained Leaves (B9) (except MLRA Water-Stained Leaves (B9) (MLRA High Water Table (A2) 1, 2, 4A, and 4B) 1, 2, 4A, and 4B) Saturation (A3) Salt Crust (B11) Drainage Patterns (B10) Water Marks (B1) Aquatic Invertebrates (B13) Dry-Season Water Table (C2) Sediment Deposits (B2) Hydrogen Sulfide Odor (C1) Saturated Visible on Aerial Imagery (C9) Drift Deposits (B3) Oxidized Rhizospheres along Living Roots (C3) Geomorphic Position (D2) Algal Mat or Crust (B4) Presence of Reduction Iron (C4) Shallow Aquitard (D3) Iron Deposits (B5) Recent Iron Reduction Tilled Soils (C6) FAC-Neutral Test (D5) Stunted or Stressed Plants (D1) (LRR A) Raised Ant Mounds (D6) (LRR A) Surface Soil Cracks (B6) Other (Explain in Remarks) Frost-Heave Hummocks (D7) Inundation Visible on Aerial Imagery (B7) Sparsely Vegetated Concave Surface (B8) Field Observations: Wetland Hydrology Present? Surface Water Present? ✓ No Depth (inches): \neg_{Yes} Water Table Present? √ No Depth (inches): Yes Yes ✓ No Yes Saturation Present? Yes ✓ No Depth (inches): (includes capillary fringe) Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available: Remarks:

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site: Newport Way Imp		y Improvements	Project	City/County:	King		Sampling Date: 5/3/201			5/3/2018
Applicant/Owner:	oplicant/Owner: City of Issaquah		f Issaquah		State	:: WA			ling Point: SF	93
Investigator(s): DBC				Section/Townsh	ip/Range: S29 T24N RC		<u> </u>			
Landform (hillslope, terr	race, etc.):	terrace		Local Relief (cor	ncave, convex, none):		concave	. :	Slope (%): <u><5</u>	5%
Subregion (LLR):	Α		_ Lat:	47.544552°	Long: <u>-122.066469</u>		Datum:	WGS 1984		
Soil Map Unit Name:	Alderwood a	and Kitsap soils, v	ery steep		_ NV	VI Classification:	NA			
Are climatic/hydrologic	conditions on t	the site typical fo	r this time of year?		✓ Yes	☐ No	(if no, explain in	n Remarks.)		
Are Vegetation Soil Hydrology			significantly distu	rbed?	Are "normal	circumstances" p	resent?	Yes No		
Are Vegetation	Soil	Hydrology	naturally problem	natic?	(if needed, ex	xplain any answe	rs in Remarks.)			
SUMMARY OF FIND Hydrophytic Vegetation		✓ Yes	1 _{N-}	T						
Hydric Soil Present? Weltand Hydrology Pres		✓ Yes ☐ ✓ Yes ☐	_ No _ No _ No	Is the sampled a Wetland?	area within a	✓ Yes	0			
,										
Remarks:										
VEGETATION - Use	scientific na	mes of plants	Absolute %	Dominant	Indicator	T				
Tree Stratum			Cover	Species?	Status	Dominance Te	st Worksheet:			
1.					-	Number of dor				(0)
2. 3.			-		-	That are OBL, F	-ACW, or FAC:		3	(A)
4.			-		-	Total Number	of Dominant			
Sapling/Shurb Strate	um		0	= Total Cover		Species A	cross All Strata:		4	(B)
1. Rubus spectabilis	•	-	15	Υ	FAC	Percent of don	ninant Species			
2. Symphoricarpos albu	s		5	Y	FACU	That are OBL, F	ACW, or FAC:		75	(A/B)
3. 4.					-	Prevalence Ind	ov Workshoots			
<i>5.</i>			-		•	_	Cover of:	Multiply by:		
			20	= Total Cover	•	OBL Species		x 1 =	0	
Herb Stratum				_		FACW Species	50	x 2 =	100	
1. Equisetum arvense			10	N	FAC	FAC Species		x 3 =	150	
2. Athyrium felix-femine 3. Phalaris arundinacea			25 50	·	FAC FACW	FACU Species UPL Species	5	x 4 =	<u>20</u> 0	
4.	'			'	FACV	Column Totals	105		270	(B)
5.								. ` '		` '
6.						Pr	evalence Index	= B/A =	2.57	
7. 8.			-		-	Hydrophytic V	egetation Indica	tors:		
9.			-				est for Hydroph		n	
10.						🗹 2 - Domina	nce Test is >509	%		
11.					-		nce Index is ≤3			
			85	= Total Cover		4 - Morpho	ological Adaptat	ions¹ (provide	supporting	data in
Woody Vine Stratur	<u>n</u>						on a separate s			
1.							d Non-Vascular			
2.				_ Tc+-1.0	-		ydrophytic Vege			
% Bare Ground	in Herb Stratu	ım	0	= Total Cover			of hydric soil unless distur		-	gy must
Remarks:						Hydrophytic	: Vegetation Pre	esent?	✓ Yes 🔲	No
						•			-	

SOIL Sampling Point: <u>SP3</u>

Depth	Matri	х	Redox Features					
(inches)	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²	Texture	Remarks
0 to 12	10 YR 2/1	100					loamy mucky mir	
12 to 18	2.5 Y 3/1	100					sandy gravelly	
¹ Type: C=Concentra				S=Covered or	Coated Sand	Grains.		re Lining, M=Matrix
Hydric Soil Indicators:	(Applicable to all LR	Rs, unless ot	herwise noted.)				Indicators for Pro	oblematic Hydric Soils ³ :
Histisol (A1)			Sandy Redox (S5)				2 cm Muck (A	A10)
Histic Epipedon (A2	2)	Ħ	Stripped Matrix (Se	5)			Red Parent M	•
Black Histic (A3)	,		Loamy Mucky Min		t MLRA 1)		=	Dard Surface (TF12)
Hydrogen Sulfide (44)	H	Loamy Gleyed Mat		·•···,		= '	n in Remarks)
Depleted Below Da		П	Depleted Matrix (F				other (Explai	nemana,
Thick Dark Surface	, ,	H	Redox Dark Surface	•			iiiuicatoi 3 Oi	nyaropnytic vegetation and
Sandy Mucky Mine	• •	Ħ	Depleted Dark Sur				wetland hydro	ology must be present,
Sandy Gleyed Matr	` '	П	Redox Depressions				unless disturb	ed or problematic.
Restrictive Layer (if pre	· '			, ,	ydric Soil Preser	nt?		
Туре					ĺ			
Depth (inches								✓ Yes No
Remarks:								
The trial has								
HYDROLOGY								
Wetland Hydrology Inc	dicators:							
Primary Indicators (mir	imum of one require	ed; check all	that apply)				Secondary Indicat	tors (2 or more required)
Surface Water (A1)			☐ Water-Stained	Leaves (B9) (ex	cept MLRA		☐ Water-Staine	d Leaves (B9) (MLRA
High Water Table (A2)		1, 2, 4A, and 4		•		1, 2, 4A, and	14B)
Saturation (A3)			Salt Crust (B11	.)			☐ Drainage Pat	terns (B10)
Water Marks (B1)			Aquatic Inverte	ebrates (B13)			Dry-Season V	Vater Table (C2)
Sediment Deposits	(B2)		Hydrogen Sulfi	de Odor (C1)			Saturated Vis	sible on Aerial Imagery (C9)
Drift Deposits (B3)			Oxidized Rhizo	spheres along Li	ving Roots (C3)		Geomorphic	Position (D2)
Algal Mat or Crust	(B4)		Presence of Re	duction Iron (C4)		Shallow Aqui	tard (D3)
Iron Deposits (B5)			Recent Iron Re	duction Tilled Sc	oils (C6)		FAC-Neutral	Test (D5)
Surface Soil Cracks	(B6)		Stunted or Stre	essed Plants (D1)	(LRR A)		Raised Ant M	lounds (D6) (LRR A)
Inundation Visible	on Aerial Imagery (B	7)	Other (Explain	in Remarks)			Frost-Heave I	Hummocks (D7)
Sparsely Vegetated	Concave Surface (B	8)						
Field Observations:				Wetla	nd Hydrology Pr	esent?		
Surface Water Present?	? Yes	✓ No	Depth (inches):					
Water Table Present?	✓ Yes	☐ No	Depth (inches):	2" bgs	.[✓ Yes No
Saturation Present?	✓ Yes	No No	Depth (inches):	0" bgs	.			
(includes capillary fring								
Describe Recorded Data	a (stream gauge, mo	nitoring well	, aerial photos, prev	ious inspections), if available:			
<u> </u>								
Remarks:								
1								

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site: Newport Way Improvement Applicant/Owner: City of Issaquah		Project City/County		King		Sampling Date: 5/3/2018			
				_ State	: WA		Sampling Point	: SP4	
Investigator(s):	DBC			Section/Townsh	ip/Range:	S29 T24N R06E			
Landform (hillslope, terr	race, etc.):	slope		Local Relief (con	cave, convex, r	none): <u>c</u>	oncave	Slope (%):	: 5%
Subregion (LLR):	Α	_	Lat:	47.544284°	Long	: <u>-122.066400°</u>	Datum:	WGS 1984	-
Soil Map Unit Name:	Alderwood a	nd Kitsap soils,	very steep		_ N	WI Classification: N	IA		-
Are climatic/hydrologic	conditions on t	he site typical fo	or this time of year?		✓ Yes	□ No (i	if no, explain	in Remarks.)	
Are Vegetation Soil Hydrology significantly			significantly distu	rbed?	Are "normal	circumstances" pres	sent?	✓ Yes] No
Are	Soil	Hydrology	naturally problem	natic?	(if needed, ex	xplain any answers i	in Remarks.)		
SUMMARY OF FIND			-lu	<u> </u>					
Hydrophytic Vegetation Hydric Soil Present? Weltand Hydrology Pres		Yes	✓ No ✓ No ✓ No	Is the sampled a Wetland?	rea within a	Yes Vo			
Remarks:									
VEGETATION - Use	scientific na	mes of plants	s.						
Tree Stratum		The or promo	Absolute %	Dominant	Indicator	Dominance Test	Worksheet:		
1. Alnus rubra			Cover 25	Species?	Status FAC	Number of domin	ant Species		
2. Acer macrophyllum			25	· Y	FACU	That are OBL, FAC		3	(A)
3.]	•		, ,
4.						Total Number of I			
Sapling/Shurb Strate	ıım		50	_ = Total Cover		Species Acros	ss All Strata:	7	(B)
1. Tsuga heterophylla	<u> </u>		10	N	FACU	Percent of domin	ant Species		
2. Abies grandis			10	N	FACU	That are OBL, FAC		43	(A/B)
3. Rubus spectabilis			20	Υ	FAC				
4. Oemleria cerasiformi	's		15	ΥΥ	FACU	Prevalence Index			
5.				Tatal Carra		Total % Cov		Multiply by:	`
Harb Ctratum			55	= Total Cover		OBL Species			-
Herb Stratum 1. Ranunculus repens			10	Υ	FAC	FACW Species FAC Species	<u>0</u>	x 2 = C x 3 = 180	_
2. Geranium robertianu	m		20	·	FACU	FACU Species		x 4 = 380	_
3. Polystichum munitun			5	N	FAC	UPL Species		x 5 = 0	_
4. Rubus parviflorus			15	Υ	FACU	Column Totals:	155	(A) 560) (B)
5.					-	4	1 1. 1.	D/A 2.54	
<i>6.</i> <i>7</i> .						Preva	alence index :	= B/A = <u>3.61</u>	_
8.						Hydrophytic Vege	tation Indica	tors:	
9.						1 - Rapid Test	for Hydroph	ytic Vegetation	
10.						2 - Dominanc			
11.						3 - Prevalence			
			50	= Total Cover		4 - Morpholo	gical Adaptat	ions ¹ (provide supporti	ing data in
Woody Vine Stratur	<u>n</u>					Remarks or or	•		
1.						5 - Wetland N			
2.						Problem Hydi			
			0	= Total Cover			-	and wetland hydro	
% Bare Ground	ın Herb Stratu	m	_			pe present, un	iess aisturi	bed or problemation	<i>.</i>
Remarks:						Hydrophytic Ve	getation Pre	sent? Yes	✓ No

SOIL Sampling Point: SP4 Depth Matrix **Redox Features** Type¹ Loc² (inches) Texture Remarks Color (moist) Color (moist) 7.5 YR 2.5/2 100 0 to 8 loam 8 to 11 2.5 Y 4/3 100 shovel refusal at 11" ¹Type: C=Concentration, D=Depletion, RM-Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix Indicators for Problematic Hydric Soils³: Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.) Histisol (A1) Sandy Redox (S5) 2 cm Muck (A10) Histic Epipedon (A2) Stripped Matrix (S6) Red Parent Material (TF2) Black Histic (A3) Loamy Mucky Mineral (F1) (except MLRA 1) ☐ Very Shallow Dard Surface (TF12) Loamy Gleyed Matrix (F2) Hydrogen Sulfide (A4) Other (Explain in Remarks) Depleted Below Dark Surface (A11) Depleted Matrix (F3) maicators or myarophytic vegetation Thick Dark Surface (A12) Redox Dark Surface (F6) and wetland hydrology must be present, Sandy Mucky Mineral (S1) Depleted Dark Surface (F7) unless disturbed or problematic. Sandy Gleyed Matrix (S4) Redox Depressions (F8) Restrictive Layer (if present): Hydric Soil Present? Type: Yes Vo Depth (inches): Remarks: **HYDROLOGY** Wetland Hydrology Indicators: Primary Indicators (minimum of one required; check all that apply) Secondary Indicators (2 or more required) Surface Water (A1) ■ Water-Stained Leaves (B9) (except MLRA Water-Stained Leaves (B9) (MLRA High Water Table (A2) 1, 2, 4A, and 4B) 1, 2, 4A, and 4B) Saturation (A3) Salt Crust (B11) Drainage Patterns (B10) Water Marks (B1) Aquatic Invertebrates (B13) Dry-Season Water Table (C2) Sediment Deposits (B2) Hydrogen Sulfide Odor (C1) Saturated Visible on Aerial Imagery (C9) Drift Deposits (B3) Oxidized Rhizospheres along Living Roots (C3) Geomorphic Position (D2) Algal Mat or Crust (B4) Presence of Reduction Iron (C4) Shallow Aquitard (D3) Iron Deposits (B5) Recent Iron Reduction Tilled Soils (C6) FAC-Neutral Test (D5) Stunted or Stressed Plants (D1) (LRR A) Raised Ant Mounds (D6) (LRR A) Surface Soil Cracks (B6) Other (Explain in Remarks) Frost-Heave Hummocks (D7) Inundation Visible on Aerial Imagery (B7) Sparsely Vegetated Concave Surface (B8) Field Observations: Wetland Hydrology Present? Surface Water Present? ✓ No Depth (inches): \neg_{Yes}

Yes Yes

✓ No

Water Table Present?

(includes capillary fringe)

Saturation Present?

Remarks:

√ No

✓ No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Yes

Yes

Depth (inches):

Depth (inches):

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site:	Newport Way Improvemen	ts Project	_City/County:	King		S	ampling Date:	5/3/2018
Applicant/Owner:	City of Issaquah			State	:: <u>WA</u>	Sa	ampling Point: SP	5
Investigator(s):	DBC		Section/Townsh	ip/Range:	S29 T24N R06E			
Landform (hillslope, terr	ace, etc.): slope		Local Relief (con	cave, convex, r	none): concave		Slope (%):	10%
Subregion (LLR):	Α	Lat:	47.543736°	Long	: <u>-122.065692°</u> Da	tum: WGS 19	984	
Soil Map Unit Name:	Alderwood and Kitsap soils,	, very steep		N	WI Classification: NA			
Are climatic/hydrologic o	conditions on the site typical	for this time of year?	•	✓ Yes	□ No (if no, e	xplain in Remar	rks.)	
Are Vegetation	Soil Hydrology	significantly distu	ırbed?	Are "normal	circumstances" present?		✓ Yes No	
Are Vegetation	Soil Hydrology	naturally problen	natic?	(if needed, ex	xplain any answers in Rem	arks.)		
SUMMARY OF FIND								
Hydrophytic Vegetation Hydric Soil Present? Weltand Hydrology Pres	Yes	No ✓ No ✓ No	Is the sampled a Wetland?	rea within a	Yes V No			
Remarks: FAC vegetation	n community qualifies as hydi	rophytic; no other pa	rameters met.					
VEGETATION - Use s	scientific names of plan	ts.						
Tree Stratum	·	Absolute %	Dominant Species 2	Indicator	Dominance Test Worksl	neet:		
1. Thuja plicata		Cover 70	Species? Y	Status FAC	Number of dominant Sp	ecies		
2. Acer macrophyllum		30	Υ	FACU	That are OBL, FACW, or	FAC:	3	(A)
3.								
4.		100	= Total Cover		Total Number of Domina		4	(B)
Sapling/Shurb Stratu	ım	100	_ = Total Cover		Species Across All St	Tala	4	(D)
1. Acer circinatum	<u> </u>	30	Υ Υ	FAC	Percent of dominant Spe	cies		
2.					That are OBL, FACW, or		75	(A/B)
3.								
<u>4. </u>					Prevalence Index Works			
5.		30	= Total Cover		Total % Cover of: OBL Species	Multiply x 1 =	y by: 0	
Herb Stratum			_		FACW Species	5 x 2 =	10	
1. Polystichum munitum	1	30	Υ	FAC	FAC Species	135 x 3 =	405	
2. Equisetum arvense		5	N	FAC	FACU Species	35 x 4 =	140	
3. Equisetum hyemale		<u>5</u> 5	N N	FACW	UPL Species	x 5 =	0	(D)
4. Rubus ursinus 5.		5	N	FACU	Column Totals:	175 (A)	555	(B)
6.					Prevalence	ndex = B/A =	3.17	
7.]	_		
8.					Hydrophytic Vegetation			
9. 10.					1 - Rapid Test for Hy 2 - Dominance Test i		lation	
11.					3 - Prevalence Index			
		45	= Total Cover		4 - Morphological Ad	daptations ¹ (pro	ovide supporting	data in
Woody Vine Stratun	<u>1</u>		_		Remarks or on a sepa			
1.					5 - Wetland Non-Vas	cular Plants ¹		
2.					Problem Hydrophyti		Explain)	
		0	= Total Cover		¹ Indicators of hydric			gy must
% Bare Ground	in Herb Stratum				be present, unless d			
Remarks: FAC vegetation	on community qualifies as h	ydrophytic			Hydrophytic Vegetation	on Present?	✓ Yes	No

SOIL Sampling Point: SP5 Depth Matrix **Redox Features** Type¹ Loc² Texture Remarks (inches) Color (moist) Color (moist) 0 to 10" 10 YR 3/2 100 loam ¹Type: C=Concentration, D=Depletion, RM-Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix Indicators for Problematic Hydric Soils³: Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.) Histisol (A1) Sandy Redox (S5) 2 cm Muck (A10) Histic Epipedon (A2) Stripped Matrix (S6) Red Parent Material (TF2) Black Histic (A3) Loamy Mucky Mineral (F1) (except MLRA 1) ☐ Very Shallow Dard Surface (TF12) Loamy Gleyed Matrix (F2) Hydrogen Sulfide (A4) Other (Explain in Remarks) Depleted Below Dark Surface (A11) Depleted Matrix (F3) maicators or myarophytic vegetation Thick Dark Surface (A12) Redox Dark Surface (F6) and wetland hydrology must be present, Sandy Mucky Mineral (S1) Depleted Dark Surface (F7) unless disturbed or problematic. Sandy Gleyed Matrix (S4) Redox Depressions (F8) Restrictive Layer (if present): Hydric Soil Present? Type: Yes Vo Depth (inches): Remarks: **HYDROLOGY** Wetland Hydrology Indicators: Primary Indicators (minimum of one required; check all that apply) Secondary Indicators (2 or more required) Surface Water (A1) □ Water-Stained Leaves (B9) (except MLRA) Water-Stained Leaves (B9) (MLRA High Water Table (A2) 1, 2, 4A, and 4B) 1, 2, 4A, and 4B) Saturation (A3) Salt Crust (B11) Drainage Patterns (B10) Water Marks (B1) Aquatic Invertebrates (B13) Dry-Season Water Table (C2) Sediment Deposits (B2) Hydrogen Sulfide Odor (C1) Saturated Visible on Aerial Imagery (C9) Drift Deposits (B3) Oxidized Rhizospheres along Living Roots (C3) Geomorphic Position (D2) Algal Mat or Crust (B4) Presence of Reduction Iron (C4) Shallow Aquitard (D3) Iron Deposits (B5) Recent Iron Reduction Tilled Soils (C6) FAC-Neutral Test (D5) Stunted or Stressed Plants (D1) (LRR A) Raised Ant Mounds (D6) (LRR A) Surface Soil Cracks (B6) Other (Explain in Remarks) Frost-Heave Hummocks (D7) Inundation Visible on Aerial Imagery (B7) Sparsely Vegetated Concave Surface (B8) Field Observations: Wetland Hydrology Present? Surface Water Present? ✓ No Depth (inches): \neg_{Yes} Water Table Present? √ No Depth (inches): Yes Yes ✓ No Yes Saturation Present? Yes ✓ No Depth (inches): (includes capillary fringe) Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available: Remarks: No hydro indicators

WETLAND DETERMINATION DATA FORM - Western Mountains, Valleys, and Coast Region

Project/Site:	Newport Way Improvem	ents Project	_City/County:	King		Sampling Date:	5/3/2018
Applicant/Owner:	City of Issaquah			State	e: WA	Sampling Point: SP	96
Investigator(s):	DBC		Section/Townsh	ip/Range:	S29 T24N R06E		
Landform (hillslope, terra	ace, etc.): slope		Local Relief (con	icave, convex, i	none): concave	Slope (%):	5%
Subregion (LLR):	Α	Lat:	47.543418°	Long	g: <u>-122.065843°</u> Datum:	WGS 1984	
Soil Map Unit Name:	Alderwood and Kitsap so	ils, very steep			IWI Classification: NA		
Are climatic/hydrologic o	conditions on the site typic	cal for this time of year?	•	✓ Yes	☐ No (if no, explain	n in Remarks.)	
Are	Soil Hydrology	significantly distu	ırbed?	Are "normal	circumstances" present?	✓ Yes No	1
Are Vegetation	Soil Hydrology	naturally probler	natic?	(if needed, e	xplain any answers in Remarks.)	1	
SUMMARY OF FIND			_				
Hydrophytic Vegetation Hydric Soil Present? Weltand Hydrology Pres	Yes	No ✓ No No	Is the sampled a Wetland?	irea within a	✓ Yes		
Remarks:							
VEGETATION - Use s	scientific names of pla				_		
Tree Stratum		Absolute % Cover	Dominant Species?	Indicator Status	Dominance Test Worksheet:		
1.					Number of dominant Species		
<i>2.</i> <i>3.</i>			-	-	That are OBL, FACW, or FAC:	3	(A)
4.					Total Number of Dominant		
		0	= Total Cover		Species Across All Strata:	3	(B)
Sapling/Shurb Stratu	<u>um</u>						
1.Rubus spectabilis		40	- Y	FAC	Percent of dominant Species	400	(4 (5)
2. Thuja plicata 3.		5	N	FAC	That are OBL, FACW, or FAC:	100	(A/B)
4.				-	Prevalence Index Worksheet:		
5.					Total % Cover of:	Multiply by:	
		45	= Total Cover		OBL Species 15	x 1 =15	
Herb Stratum		2-	.,		FACW Species 30		
Equisetum arvense Phalaris arundinacea		25 30	- Y Y	FAC FACW	FAC Species 85 FACU Species 0		
3. Athyrium felix-femina	1	15	- <u>'</u>	FAC		x5= 0	
4. Lysichiton americanus		15	N	OBL		(A) 330	(B)
5.]	· . ——	
<i>6.</i> <i>7.</i>				_	Prevalence Index	z = B/A = 2.54	
<i>8.</i>				-	Hydrophytic Vegetation Indic	ators:	
9.				_	1 - Rapid Test for Hydrop		
10.					✓ 2 - Dominance Test is >50	0%	
11.					3 - Prevalence Index is ≤3		
		85	= Total Cover		4 - Morphological Adapta	tions ¹ (provide supporting	data in
Woody Vine Stratun	<u>n</u>				Remarks or on a separate		
1.					5 - Wetland Non-Vascular		
2.			-		Problem Hydrophytic Veg		
		0	= Total Cover		¹ Indicators of hydric soil		gy must
% Bare Ground	in Herb Stratum				be present, unless distu	rbed or problematic.	
Remarks:					Hydrophytic Vegetation Pro	esent?	No

SOIL Sampling Point: SP6 Matrix **Redox Features** Depth Type¹ Loc² Texture Remarks (inches) Color (moist) % Color (moist) 10 YR 3/2 0 to 10" 100 loam 10 to 16" 10 YR 3/2 90 10 YR 4/6 10 loam ¹Type: C=Concentration, D=Depletion, RM-Reduced Matrix, CS=Covered or Coated Sand Grains. ²Location: PL=Pore Lining, M=Matrix Indicators for Problematic Hydric Soils³: Hydric Soil Indicators: (Applicable to all LRRs, unless otherwise noted.) Histisol (A1) Sandy Redox (S5) 2 cm Muck (A10) Histic Epipedon (A2) Stripped Matrix (S6) Red Parent Material (TF2) Black Histic (A3) Loamy Mucky Mineral (F1) (except MLRA 1) ☐ Very Shallow Dard Surface (TF12) Loamy Gleyed Matrix (F2) Hydrogen Sulfide (A4) Other (Explain in Remarks) Depleted Below Dark Surface (A11) Depleted Matrix (F3) maicators or myarophytic vegetation Thick Dark Surface (A12) Redox Dark Surface (F6) and wetland hydrology must be present, Sandy Mucky Mineral (S1) Depleted Dark Surface (F7) unless disturbed or problematic. Sandy Gleyed Matrix (S4) Redox Depressions (F8) Restrictive Layer (if present): Hydric Soil Present? Type: Yes Vo Depth (inches): Remarks: redox layer too deep to qualify as indicator F6; plot located at wetland margin - assumed shallower groundwater saturation at interior. **HYDROLOGY** Wetland Hydrology Indicators: Primary Indicators (minimum of one required; check all that apply) Secondary Indicators (2 or more required) Surface Water (A1) □ Water-Stained Leaves (B9) (except MLRA) Water-Stained Leaves (B9) (MLRA ✓ High Water Table (A2) 1, 2, 4A, and 4B) 1, 2, 4A, and 4B) ✓ Saturation (A3) Salt Crust (B11) Drainage Patterns (B10) Water Marks (B1) Aquatic Invertebrates (B13) Dry-Season Water Table (C2) Sediment Deposits (B2) Hydrogen Sulfide Odor (C1) Saturated Visible on Aerial Imagery (C9) Drift Deposits (B3) Oxidized Rhizospheres along Living Roots (C3) Geomorphic Position (D2) Algal Mat or Crust (B4) Presence of Reduction Iron (C4) Shallow Aquitard (D3) Iron Deposits (B5) Recent Iron Reduction Tilled Soils (C6) FAC-Neutral Test (D5) Stunted or Stressed Plants (D1) (LRR A) Raised Ant Mounds (D6) (LRR A) Surface Soil Cracks (B6) Other (Explain in Remarks) Frost-Heave Hummocks (D7) Inundation Visible on Aerial Imagery (B7) Sparsely Vegetated Concave Surface (B8) Field Observations: Wetland Hydrology Present? Surface Water Present? ✓ No Depth (inches): \neg_{Yes}

✓ Yes

ΠNo

Depth (inches):

Depth (inches):

4" bgs 0" bgs

✓ Yes

✓ Yes

No

No

Describe Recorded Data (stream gauge, monitoring well, aerial photos, previous inspections), if available:

Water Table Present?

(includes capillary fringe)

Saturation Present?

Remarks:

APPENDIX DWetland Rating Form

RATING SUMMARY – Western Washington

Name of wetland (or ID #): Wetland A Date of site visit: 10/30/17 Rated by E. Duncanson Trained by Ecology? X Yes ____No Date of training: June 2017

HGM Class used for rating: Riverine Wetland has multiple HGM classes?_ Y _x_ N

NOTE: Form is not complete without the figures requested (figures can be combined).

Source of base aerial photo/map - <u>ESRI ArcGIS</u>

OVERALL WETLAND CATEGORY __II__ (based on functions__ or special characteristics__)

1. Category of wetland based on FUNCTIONS

Category I – Total score = 23 - 27

X Category II – Total score = 20 - 22

Category III – Total score = 16 - 19

Category IV – Total score = 9 - 15

FUNCTION	Improving Water Quality		H	ydrol	ogic		Habita	it		
					Circle	the ap	propi	riate ra	tings	
Site Potential	Н	M	L	H) M	L	Н	М	(L)	
Landscape Potential	Н	M	L	H) M	L	Н	M	L	
Value (Н	М	L (Н) M	L	H	М	L	TOTAL
Score Based on Ratings		7			9			6		22

Score for each function based on three ratings (order of ratings is not *important)* 9 = H,H,H8 = H, H, M7 = H,H,L 7 = H, M, M6 = H,M,L6 = M,M,M5 = H,L,L5 = M,M,L4 = M, L, L3 = L, L, L

2. Category based on SPECIAL CHARACTERISTICS of wetland

CHARACTERISTIC	CATEGORY		
Estuarine	I II		
Wetland of High Conservation Value	I		
Bog	I		
Mature Forest	I		
Old Growth Forest	I		
Coastal Lagoon	I II		
Interdunal	I II	III IV	
None of the above	NA		

Maps and figures required to answer questions correctly for Western Washington

Depressional Wetlands

Map of:	To answer questions:	Figure #
Cowardin plant classes	D 1.3, H 1.1, H 1.4	
Hydroperiods	D 1.4, H 1.2	
Location of outlet (can be added to map of hydroperiods)	D 1.1, D 4.1	
Boundary of area within 150 ft of the wetland (can be added to another figure)	D 2.2, D 5.2	
Map of the contributing basin	D 4.3, D 5.3	
1 km Polygon: Area that extends 1 km from entire wetland edge - including	H 2.1, H 2.2, H 2.3	
polygons for accessible habitat and undisturbed habitat		
Screen capture of map of 303(d) listed waters in basin (from Ecology website)	D 3.1, D 3.2	
Screen capture of list of TMDLs for WRIA in which unit is found (from web)	D 3.3	

Riverine Wetlands

Map of:	To answer questions:	Figure #
Cowardin plant classes	H 1.1, H 1.4	1
Hydroperiods	H 1.2	1
Ponded depressions	R 1.1	1
Boundary of area within 150 ft of the wetland (can be added to another figure)	R 2.4	1
Plant cover of trees, shrubs, and herbaceous plants	R 1.2, R 4.2	1
Width of unit vs. width of stream (can be added to another figure)	R 4.1	1
Map of the contributing basin	R 2.2, R 2.3, R 5.2	2
1 km Polygon: Area that extends 1 km from entire wetland edge - including	H 2.1, H 2.2, H 2.3	3
polygons for accessible habitat and undisturbed habitat		3
Screen capture of map of 303(d) listed waters in basin (from Ecology website)	R 3.1	4
Screen capture of list of TMDLs for WRIA in which unit is found (from web)	R 3.2, R 3.3	5

Lake Fringe Wetlands

Map of:	To answer questions:	Figure #
Cowardin plant classes	L 1.1, L 4.1, H 1.1, H 1.4	
Plant cover of trees, shrubs, and herbaceous plants	L 1.2	
Boundary of area within 150 ft of the wetland (can be added to another figure)	L 2.2	
1 km Polygon: Area that extends 1 km from entire wetland edge - including	H 2.1, H 2.2, H 2.3	
polygons for accessible habitat and undisturbed habitat		
Screen capture of map of 303(d) listed waters in basin (from Ecology website)	L 3.1, L 3.2	
Screen capture of list of TMDLs for WRIA in which unit is found (from web)	L 3.3	

Slope Wetlands

Map of:	To answer questions:	Figure #
Cowardin plant classes	H 1.1, H 1.4	
Hydroperiods	H 1.2	
Plant cover of dense trees, shrubs, and herbaceous plants	S 1.3	
Plant cover of dense, rigid trees, shrubs, and herbaceous plants	S 4.1	
(can be added to figure above)		
Boundary of 150 ft buffer (can be added to another figure)	S 2.1, S 5.1	
1 km Polygon: Area that extends 1 km from entire wetland edge - including	H 2.1, H 2.2, H 2.3	
polygons for accessible habitat and undisturbed habitat		
Screen capture of map of 303(d) listed waters in basin (from Ecology website)	S 3.1, S 3.2	
Screen capture of list of TMDLs for WRIA in which unit is found (from web)	S 3.3	

HGM Classification of Wetlands in Western Washington

For questions 1-7, the criteria described must apply to the entire unit being rated.

If the hydrologic criteria listed in each question do not apply to the entire unit being rated, you probably have a unit with multiple HGM classes. In this case, identify which hydrologic criteria in questions 1-7 apply, and go to Question 8.

1. Are the water levels in the entire unit usually controlled by tides except during floods?

NO - go to 2

YES – the wetland class is **Tidal Fringe** – go to 1.1

1.1 Is the salinity of the water during periods of annual low flow below 0.5 ppt (parts per thousand)?

NO - Saltwater Tidal Fringe (Estuarine)

YES - Freshwater Tidal Fringe

If your wetland can be classified as a Freshwater Tidal Fringe use the forms for **Riverine** wetlands. If it is Saltwater Tidal Fringe it is an **Estuarine** wetland and is not scored. This method **cannot** be used to score functions for estuarine wetlands.

2. The entire wetland unit is flat and precipitation is the only source (>90%) of water to it. Groundwater and surface water runoff are NOT sources of water to the unit.

NO - go to 3

YES - The wetland class is Flats

If your wetland can be classified as a Flats wetland, use the form for **Depressional** wetlands.

- 3. Does the entire wetland unit **meet all** of the following criteria?
 - __The vegetated part of the wetland is on the shores of a body of permanent open water (without any plants on the surface at any time of the year) at least 20 ac (8 ha) in size;
 - __At least 30% of the open water area is deeper than 6.6 ft (2 m).

NO - go to 4

YES - The wetland class is **Lake Fringe** (Lacustrine Fringe)

- 4. Does the entire wetland unit **meet all** of the following criteria?
 - __The wetland is on a slope (*slope can be very gradual*),
 - __The water flows through the wetland in one direction (unidirectional) and usually comes from seeps. It may flow subsurface, as sheetflow, or in a swale without distinct banks,
 - __The water leaves the wetland **without being impounded**.

NO - go to 5

YES - The wetland class is **Slope**

NOTE: Surface water does not pond in these type of wetlands except occasionally in very small and shallow depressions or behind hummocks (depressions are usually <3 ft diameter and less than 1 ft deep).

- 5. Does the entire wetland unit **meet all** of the following criteria?
 - _x_The unit is in a valley, or stream channel, where it gets inundated by overbank flooding from that stream or river,
 - _x_The overbank flooding occurs at least once every 2 years.

Wetland name or number A

NO – go to 6

YES - The wetland class is Riverine

NOTE: The Riverine unit can contain depressions that are filled with water when the river is not flooding

6. Is the entire wetland unit in a topographic depression in which water ponds, or is saturated to the surface, at some time during the year? *This means that any outlet, if present, is higher than the interior of the wetland.*

NO - go to 7

YES – The wetland class is **Depressional**

7. Is the entire wetland unit located in a very flat area with no obvious depression and no overbank flooding? The unit does not pond surface water more than a few inches. The unit seems to be maintained by high groundwater in the area. The wetland may be ditched, but has no obvious natural outlet.

NO - go to 8

YES - The wetland class is **Depressional**

8. Your wetland unit seems to be difficult to classify and probably contains several different HGM classes. For example, seeps at the base of a slope may grade into a riverine floodplain, or a small stream within a Depressional wetland has a zone of flooding along its sides. GO BACK AND IDENTIFY WHICH OF THE HYDROLOGIC REGIMES DESCRIBED IN QUESTIONS 1-7 APPLY TO DIFFERENT AREAS IN THE UNIT (make a rough sketch to help you decide). Use the following table to identify the appropriate class to use for the rating system if you have several HGM classes present within the wetland unit being scored.

NOTE: Use this table only if the class that is recommended in the second column represents 10% or more of the total area of the wetland unit being rated. If the area of the HGM class listed in column 2 is less than 10% of the unit; classify the wetland using the class that represents more than 90% of the total area.

HGM classes within the wetland unit	HGM class to
being rated	use in rating
Slope + Riverine	Riverine
Slope + Depressional	Depressional
Slope + Lake Fringe	Lake Fringe
Depressional + Riverine along stream	Depressional
within boundary of depression	
Depressional + Lake Fringe	Depressional
Riverine + Lake Fringe	Riverine
Salt Water Tidal Fringe and any other	Treat as
class of freshwater wetland	ESTUARINE

If you are still unable to determine which of the above criteria apply to your wetland, or if you have **more than 2 HGM classes** within a wetland boundary, classify the wetland as Depressional for the rating.

- Vectoria name of number	
RIVERINE AND FRESHWATER TIDAL FRINGE WETLANDS	
Water Quality Functions - Indicators that the site functions to improve water quality	
R 1.0. Does the site have the potential to improve water quality?	
R 1.1. Area of surface depressions within the Riverine wetland that can trap sediments during a flooding event:	
Depressions cover $> \frac{3}{4}$ area of wetland points = 8	
Depressions cover > ½ area of wetland points = 4	
Depressions present but cover < ½ area of wetland points = 2	2
No depressions present points = 0	
R 1.2. Structure of plants in the wetland (areas with >90% cover at person height, not Cowardin classes)	
Trees or shrubs $> \frac{2}{3}$ area of the wetland points = 8	
Trees or shrubs $> \frac{1}{3}$ area of the wetland points = 6	
Herbaceous plants (> 6 in high) > $^{2}/_{3}$ area of the wetland points = 6	8
Herbaceous plants (> 6 in high) > $\frac{1}{3}$ area of the wetland points = 3	
Trees, shrubs, and ungrazed herbaceous $< \frac{1}{3}$ area of the wetland points = 0	
Total for R 1 Add the points in the boxes above	10
Rating of Site Potential If score is: 12-16 = H	the first page
R 2.0. Does the landscape have the potential to support the water quality function of the site?	-
R 2.1. Is the wetland within an incorporated city or within its UGA? Yes = 2 No = 0	2
R 2.2. Does the contributing basin to the wetland include a UGA or incorporated area? Yes = 1 No = 0	1
R 2.3. Does at least 10% of the contributing basin contain tilled fields, pastures, or forests that have been clearcut	
within the last 5 years? Yes = 1 No = 0	0
R 2.4. Is $> 10\%$ of the area within 150 ft of the wetland in land uses that generate pollutants? Yes = 1 No = 0	1
R 2.5. Are there other sources of pollutants coming into the wetland that are not listed in questions R 2.1-R 2.4	0
Other sources Yes = 1 No = 0	
Total for R 2 Add the points in the boxes above	4
Rating of Landscape Potential If score is: X 3-6 = H 1 or 2 = M 0 = L Record the rating on the score is: X 3-6 = H 1 or 2 = M 1 or 2 = M 2 or 3 or	he first page
R 3.0. Is the water quality improvement provided by the site valuable to society?	
R 3.1. Is the wetland along a stream or river that is on the 303(d) list or on a tributary that drains to one within 1 mi?	
	1
Yes = 1 No = 0	
R 3.2. Is the wetland along a stream or river that has TMDL limits for nutrients, toxics, or pathogens?	1

YES if there is a TMDL for the drainage in which the unit is found)Yes = 2 No = 0Total for R 3Add the points in the boxes above4Rating of Value If score is: $\frac{X}{2}$ -4 = H $\frac{1}{2}$ = M $\frac{0}{2}$ = LRecord the rating on the first page

R 3.3. Has the site been identified in a watershed or local plan as important for maintaining water quality? (answer

Yes = 1 No = 0

1

2

RIVERINE AND FRESHWATER TIDAL FRING	E WETLANDS	
Hydrologic Functions - Indicators that site functions to reduce	flooding and stream erosio	n
R 4.0. Does the site have the potential to reduce flooding and erosion?		
R 4.1. Characteristics of the overbank storage the wetland provides:		
Estimate the average width of the wetland perpendicular to the direction of the f stream or river channel (distance between banks). Calculate the ratio: (average width of stream between banks).	= = = = = = = = = = = = = = = = = = = =	
If the ratio is more than 20	points = 9	
If the ratio is 10-20	points = 6	6
If the ratio is 5-<10	points = 4	
If the ratio is 1-<5	points = 2	
If the ratio is < 1	points = 1	
R 4.2. Characteristics of plants that slow down water velocities during floods: <i>Treat largery shrub. Choose the points appropriate for the best description (polygons need to height. These are NOT Cowardin classes).</i>	=	
Forest or shrub for $>^1/_3$ area OR emergent plants $>^2/_3$ area	points = 7	7
Forest or shrub for $> \frac{1}{10}$ area OR emergent plants $> \frac{1}{3}$ area	points = 4	
Plants do not meet above criteria	points = 0	
Total for R 4 Add t	the points in the boxes above	13
R 5.0. Does the landscape have the potential to support the hydrologic function R 5.1. Is the stream or river adjacent to the wetland downcut?	Yes = 0 No = 1	1
R 5.1. Is the stream or river adjacent to the wetland downcut?	Yes = U NO = 1	1
R 5.2. Does the up-gradient watershed include a UGA or incorporated area?	Yes = 1 No = 0	1
R 5.3. Is the up-gradient stream or river controlled by dams?	Yes = 0 No = 1	1
Total for R 5 Add t	he points in the boxes above	3
Rating of Landscape Potential If score is: X3 = H 1 or 2 = M 0 = L	Record the rating on t	he first page
R 6.0. Are the hydrologic functions provided by the site valuable to society?		
R 6.1. Distance to the nearest areas downstream that have flooding problems? Choose the description that best fits the site. The sub-basin immediately down-gradient of the wetland has flooding problems human or natural resources (e.g., houses or salmon redds) Surface flooding problems are in a sub-basin farther down-gradient	points = 2 points = 1	1
No flooding problems anywhere downstream	points = 0	
R 6.2. Has the site been identified as important for flood storage or flood conveyance in	n a regional flood control plan? Yes = 2 No = 0	2
Total for R 6 Add t	the points in the boxes above	3
Rating of Value If score is: x 2-4 = H1 = M0 = L	Record the rating on t	he first page

These questions apply to wetlands of all HGM classes. **HABITAT FUNCTIONS** - Indicators that site functions to provide important habitat H 1.0. Does the site have the potential to provide habitat? H 1.1. Structure of plant community: Indicators are Cowardin classes and strata within the Forested class. Check the Cowardin plant classes in the wetland. Up to 10 patches may be combined for each class to meet the threshold of ¼ ac or more than 10% of the unit if it is smaller than 2.5 ac. Add the number of structures checked. Aquatic bed 4 structures or more: points = 4 1 3 structures: points = 2 ____Emergent 2 structures: points = 1 Scrub-shrub (areas where shrubs have > 30% cover) 1 structure: points = 0 _x__Forested (areas where trees have > 30% cover) If the unit has a Forested class, check if: X The Forested class has 3 out of 5 strata (canopy, sub-canopy, shrubs, herbaceous, moss/ground-cover) that each cover 20% within the Forested polygon H 1.2. Hydroperiods Check the types of water regimes (hydroperiods) present within the wetland. The water regime has to cover more than 10% of the wetland or ¼ ac to count (see text for descriptions of hydroperiods). Permanently flooded or inundated 4 or more types present: points = 3 x Seasonally flooded or inundated 3 types present: points = 2 2 Occasionally flooded or inundated 2 types present: points = 1 ^X Saturated only 1 type present: points = 0 X Permanently flowing stream or river in, or adjacent to, the wetland Seasonally flowing stream in, or adjacent to, the wetland 2 points Lake Fringe wetland 2 points Freshwater tidal wetland H 1.3. Richness of plant species Count the number of plant species in the wetland that cover at least 10 ft². Different patches of the same species can be combined to meet the size threshold and you do not have to name the species. Do not include Eurasian milfoil, reed canarygrass, purple loosestrife, Canadian thistle 1 If you counted: > 19 species points = 2 5 - 19 species points = 1 points = 0 < 5 species H 1.4. Interspersion of habitats Decide from the diagrams below whether interspersion among Cowardin plants classes (described in H 1.1), or the classes and unvegetated areas (can include open water or mudflats) is high, moderate, low, or none. If you have four or more plant classes or three classes and open water, the rating is always high. 0 None = 0 points Moderate = 2 points Low = 1 point All three diagrams in this row are **HIGH** = 3points

H 1.5. Special habitat features:	
Check the habitat features that are present in the wetland. <i>The number of checks is the number of points</i> . _X_Large, downed, woody debris within the wetland (> 4 in diameter and 6 ft long).	
$\frac{x}{x}$ Standing snags (dbh > 4 in) within the wetland	
Undercut banks are present for at least 6.6 ft (2 m) and/or overhanging plants extends at least 3.3 ft (1 m)	
over a stream (or ditch) in, or contiguous with the wetland, for at least 3.3 ft (10 m)	
Stable steep banks of fine material that might be used by beaver or muskrat for denning (> 30 degree	2
slope) OR signs of recent beaver activity are present (cut shrubs or trees that have not yet weathered	
where wood is exposed)	
At least ¼ ac of thin-stemmed persistent plants or woody branches are present in areas that are	
permanently or seasonally inundated (structures for egg-laying by amphibians)	
Invasive plants cover less than 25% of the wetland area in every stratum of plants (see H 1.1 for list of	
Total for H 1 Add the points in the boxes above	6
Rating of Site Potential If score is:15-18 = H7-14 = MX 0-6 = L Record the rating on	the first page
H 2.0. Does the landscape have the potential to support the habitat functions of the site?	
H 2.1. Accessible habitat (include only habitat that directly abuts wetland unit).	
Calculate: % undisturbed habitat $\frac{30}{2}$ + [(% moderate and low intensity land uses)/2] $\frac{5}{2}$ = $\frac{35}{2}$ %	
If total accessible habitat is:	
$> \frac{1}{3}$ (33.3%) of 1 km Polygon points = 3	1
20-33% of 1 km Polygon points = 2	
10-19% of 1 km Polygon points = 1	
< 10% of 1 km Polygon points = 0	
H 2.2. Undisturbed habitat in 1 km Polygon around the wetland.	
Calculate: % undisturbed habitat 30 + [(% moderate and low intensity land uses)/2] 5 = 35 %	
Undisturbed habitat > 50% of Polygon points = 3	
Undisturbed habitat 10-50% and in 1-3 patches points = 2	1
Undisturbed habitat 10-50% and > 3 patches points = 1	
Undisturbed habitat < 10% of 1 km Polygon points = 0	
H 2.3. Land use intensity in 1 km Polygon: If	_
> 50% of 1 km Polygon is high intensity land use points = (-2)	0
≤ 50% of 1 km Polygon is high intensity points = 0	
Total for H 2 Add the points in the boxes above	2
Rating of Landscape Potential If score is: $4-6 = H$ $x_1-3 = M$ $x_1-3 = M$ Record the rating on the score is: $x_1-x_2 = M$	he first page
H 3.0. Is the habitat provided by the site valuable to society?	-
H 3.1. Does the site provide habitat for species valued in laws, regulations, or policies? <i>Choose only the highest score</i>	
that applies to the wetland being rated.	
Site meets ANY of the following criteria: points = 2	
X It has 3 or more priority habitats within 100 m (see next page)	
The provides habitat for Threatened or Endangered species (any plant or animal on the state or federal lists)	2
— It is mapped as a location for an individual WDFW priority species	_
 It is a Wetland of High Conservation Value as determined by the Department of Natural Resources It has been categorized as an important habitat site in a local or regional comprehensive plan, in a 	
Shoreline Master Plan, or in a watershed plan	
Site has 1 or 2 priority habitats (listed on next page) within 100 m	
Site does not meet any of the criteria above points = 0	
Rating of Value If score is: $\frac{X}{2}$ = H $\frac{1}{1}$ = M $\frac{0}{1}$ = L Record the rating on	the first page

WDFW Priority Habitats

<u>Priority habitats listed by WDFW</u> (see complete descriptions of WDFW priority habitats, and the counties in which they can be found, in: Washington Department of Fish and Wildlife. 2008. Priority Habitat and Species List. Olympia, Washington. 177 pp. http://wdfw.wa.gov/publications/00165/wdfw00165.pdf or access the list from here: http://wdfw.wa.gov/conservation/phs/list/)

Count how many of the following priority habitats are within 330 ft (100 m) of the wetland unit: *NOTE:* This question is independent of the land use between the wetland unit and the priority habitat.

- **Aspen Stands:** Pure or mixed stands of aspen greater than 1 ac (0.4 ha).
- **Biodiversity Areas and Corridors**: Areas of habitat that are relatively important to various species of native fish and wildlife (*full descriptions in WDFW PHS report*).
- **Herbaceous Balds:** Variable size patches of grass and forbs on shallow soils over bedrock.
- Old-growth/Mature forests: Old-growth west of Cascade crest Stands of at least 2 tree species, forming a multi-layered canopy with occasional small openings; with at least 8 trees/ac (20 trees/ha) > 32 in (81 cm) dbh or > 200 years of age. Mature forests Stands with average diameters exceeding 21 in (53 cm) dbh; crown cover may be less than 100%; decay, decadence, numbers of snags, and quantity of large downed material is generally less than that found in old-growth; 80-200 years old west of the Cascade crest.
- **Oregon White Oak:** Woodland stands of pure oak or oak/conifer associations where canopy coverage of the oak component is important (*full descriptions in WDFW PHS report p. 158 see web link above*).
- Riparian: The area adjacent to aquatic systems with flowing water that contains elements of both aquatic and terrestrial ecosystems which mutually influence each other.
- **Westside Prairies:** Herbaceous, non-forested plant communities that can either take the form of a dry prairie or a wet prairie (*full descriptions in WDFW PHS report p. 161 see web link above*).
- Instream: The combination of physical, biological, and chemical processes and conditions that interact to provide functional life history requirements for instream fish and wildlife resources.
- **Nearshore**: Relatively undisturbed nearshore habitats. These include Coastal Nearshore, Open Coast Nearshore, and Puget Sound Nearshore. (*full descriptions of habitats and the definition of relatively undisturbed are in WDFW report see web link on previous page*).
- **Caves:** A naturally occurring cavity, recess, void, or system of interconnected passages under the earth in soils, rock, ice, or other geological formations and is large enough to contain a human.
- **Cliffs:** Greater than 25 ft (7.6 m) high and occurring below 5000 ft elevation.
- **Talus:** Homogenous areas of rock rubble ranging in average size 0.5 6.5 ft (0.15 2.0 m), composed of basalt, andesite, and/or sedimentary rock, including riprap slides and mine tailings. May be associated with cliffs.
- Snags and Logs: Trees are considered snags if they are dead or dying and exhibit sufficient decay characteristics to enable cavity excavation/use by wildlife. Priority snags have a diameter at breast height of > 20 in (51 cm) in western Washington and are > 6.5 ft (2 m) in height. Priority logs are > 12 in (30 cm) in diameter at the largest end, and > 20 ft (6 m) long.

Note: All vegetated wetlands are by definition a priority habitat but are not included in this list because they are addressed elsewhere.


CATEGORIZATION BASED ON SPECIAL CHARACTERISTICS

CATEGORIZATION BASED ON SPECIAL CHARACTERISTICS	C-1
Wetland Type	Category
Check off any criteria that apply to the wetland. Circle the category when the appropriate criteria are met.	
SC 1.0. Estuarine wetlands	
Does the wetland meet the following criteria for Estuarine wetlands?	
— The dominant water regime is tidal,	
— Vegetated, and	
— With a salinity greater than 0.5 ppt Yes –Go to SC 1.1 No= Not an estuarine wetland	
SC 1.1. Is the wetland within a National Wildlife Refuge, National Park, National Estuary Reserve, Natural Area Preserve, State Park or Educational, Environmental, or Scientific Reserve designated under WAC 332-30-151? Yes = Category I No-Go to SC 1.2	Cat. I
SC 1.2. Is the wetland unit at least 1 ac in size and meets at least two of the following three conditions? — The wetland is relatively undisturbed (has no diking, ditching, filling, cultivation, grazing, and has less than 10% cover of non-native plant species. (If non-native species are <i>Spartina</i> , see page 25) — At least ¾ of the landward edge of the wetland has a 100 ft buffer of shrub, forest, or un-grazed or un-	Cat. I
mowed grassland.	
— The wetland has at least two of the following features: tidal channels, depressions with open water, or contiguous freshwater wetlands. Yes = Category I No = Category II	Cat. II
SC 2.0. Wetlands of High Conservation Value (WHCV)	
SC 2.1. Has the WA Department of Natural Resources updated their website to include the list of Wetlands of High	
Conservation Value? Yes – Go to SC 2.2 No – Go to SC 2.3	Cat. I
SC 2.2. Is the wetland listed on the WDNR database as a Wetland of High Conservation Value? Yes = Category I No = Not a WHCV	
SC 2.3. Is the wetland in a Section/Township/Range that contains a Natural Heritage wetland?	
http://www1.dnr.wa.gov/nhp/refdesk/datasearch/wnhpwetlands.pdf	
Yes – Contact WNHP/WDNR and go to SC 2.4 No = Not a WHCV	
SC 2.4. Has WDNR identified the wetland within the S/T/R as a Wetland of High Conservation Value and listed it on	
their website? Yes = Category I (No)= Not a WHCV	
SC 3.0. Bogs	
Does the wetland (or any part of the unit) meet both the criteria for soils and vegetation in bogs? Use the key	
below. If you answer YES you will still need to rate the wetland based on its functions.	
SC 3.1. Does an area within the wetland unit have organic soil horizons, either peats or mucks, that compose 16 in or	
more of the first 32 in of the soil profile? Yes – Go to SC 3.3 (No) – Go to SC 3.2	
SC 3.2. Does an area within the wetland unit have organic soils, either peats or mucks, that are less than 16 in deep	
over bedrock, or an impermeable hardpan such as clay or volcanic ash, or that are floating on top of a lake or pond? Yes – Go to SC 3.3 No = Is not a bog	
SC 3.3. Does an area with peats or mucks have more than 70% cover of mosses at ground level, AND at least a 30%	
cover of plant species listed in Table 4? Yes = Is a Category I bog No - Go to SC 3.4	
NOTE: If you are uncertain about the extent of mosses in the understory, you may substitute that criterion by	
measuring the pH of the water that seeps into a hole dug at least 16 in deep. If the pH is less than 5.0 and the	Cat
plant species in Table 4 are present, the wetland is a bog.	Cat. I
SC 3.4. Is an area with peats or mucks forested (> 30% cover) with Sitka spruce, subalpine fir, western red cedar,	
western hemlock, lodgepole pine, quaking aspen, Engelmann spruce, or western white pine, AND any of the	
species (or combination of species) listed in Table 4 provide more than 30% of the cover under the canopy?	
Yes = Is a Category I bog No = Is not a bog	


Does the wetland have at least <u>1 contiguous acre</u> of forest that meets one of these criteria for the WA Department of Fish and Wildlife's forests as priority habitats? <i>If you answer YES you will still need to rate the wetland based on its functions.</i>	
 Old-growth forests (west of Cascade crest): Stands of at least two tree species, forming a multi-layered canopy with occasional small openings; with at least 8 trees/ac (20 trees/ha) that are at least 200 years of age OR have a diameter at breast height (dbh) of 32 in (81 cm) or more. Mature forests (west of the Cascade Crest): Stands where the largest trees are 80- 200 years old OR the species that make up the canopy have an average diameter (dbh) exceeding 21 in (53 cm). 	
Yes = Category I No = Not a forested wetland for this section	Cat. I
SC 5.0. Wetlands in Coastal Lagoons	
Does the wetland meet all of the following criteria of a wetland in a coastal lagoon? — The wetland lies in a depression adjacent to marine waters that is wholly or partially separated from marine waters by sandbanks, gravel banks, shingle, or, less frequently, rocks	
— The lagoon in which the wetland is located contains ponded water that is saline or brackish (> 0.5 ppt) during most of the year in at least a portion of the lagoon (needs to be measured near the bottom) Yes – Go to SC 5.1 No = Not a wetland in a coastal lagoon	Cat. I
GC 5.1. Does the wetland meet all of the following three conditions? — The wetland is relatively undisturbed (has no diking, ditching, filling, cultivation, grazing), and has less than 20% cover of aggressive, opportunistic plant species (see list of species on p. 100).	Cat. II
— At least ¾ of the landward edge of the wetland has a 100 ft buffer of shrub, forest, or un-grazed or unmowed grassland.	
— The wetland is larger than $^{1}/_{10}$ ac (4350 ft ²) Yes = Category I No = Category II	
C 6.0. Interdunal Wetlands	
Is the wetland west of the 1889 line (also called the Western Boundary of Upland Ownership or WBUO)? If you answer yes you will still need to rate the wetland based on its habitat functions.	
In practical terms that means the following geographic areas:	
Long Beach Peninsula: Lands west of SR 103	
— Grayland-Westport: Lands west of SR 105	Cat I
 Ocean Shores-Copalis: Lands west of SR 115 and SR 109 Yes – Go to SC 6.1 No= not an interdunal wetland for rating	
SC 6.1. Is the wetland 1 ac or larger and scores an 8 or 9 for the habitat functions on the form (rates H,H,H or H,H,M	Cat. II
for the three aspects of function)? Yes = Category I No – Go to SC 6.2	
C 6.2. Is the wetland 1 ac or larger, or is it in a mosaic of wetlands that is 1 ac or larger?	.
Yes = Category II No – Go to SC 6.3	Cat. III
IC 6.3. Is the unit between 0.1 and 1 ac, or is it in a mosaic of wetlands that is between 0.1 and 1 ac?	
Yes = Category III No = Category IV	Cat. IV

Wetland name or number A

This page left blank intentionally




Legend



Forest

Emergent





Notes:

The locations of all features shown are approximate.
 This drawing is for information purposes. It is intended to assist in showing features discussed in an attached document. GeoEngineers, Inc. cannot guarantee the accuracy and content of electronic files. The master file.

document. GeoEngineers, Inc. cannot guarantee the accuracy and content of electronic files. The master file is stored by GeoEngineers, Inc. and will serve as the official record of this communication.

Data Source: King County GIS, ESRI

Projection: NAD 1983 StatePlane Washington North FIPS 4601 Feet

Boundaries and Characteristics of Wetland A

Newport Way Improvements Project Issaquah, Washington



Figure D-1




- 1. The locations of all features shown are approximate.
- This drawing is for information purposes. It is intended to
 assist in showing features discussed in an attached document.
 GeoEngineers, Inc. cannot guarantee the accuracy and content
 of electronic files. The master file is stored by GeoEngineers,
 Inc. and will serve as the official record of this communication.

Data Source: King County GIS, ESRI

Projection: NAD 1983 StatePlane Washington North FIPS 4601 Feet







Landscape Characteristics of Wetland A

Newport Way Improvements Project Issaquah, Washington



Figure D-2




Figure D-3. Contributing Basin





	"The JO3(d) List contains only Category 5 Listings."								
View	ListingID	Assessment Unit ID	Medium	Parameter	Current Category	Waterbody Name	WHIA	WO Improvement Project	WQ Atlas Map Link
View	4808	17110012000221	Water	Temperature	.5	TIBBETTS CREEK	8 - Cedar-Sammamish		4808
(View.)	12683	17110012000221	Water	Dissolved Oxygen	-5	TIBBETTS CREEK	8 - Cedar-Sammamish		12683
[View.]	15778	17110012000224	Water	Dissolved Oxygen	5	TIBBETTS CREEK	8 - Cedar-Sammamish		15778
[View.]	15781	17110012000224	Water	Temperature	.5	TIBBETTS CREEK	8 - Cedar-Sammamish		15781
(View)	70112	17110012000224	Other	Bioassessment	5	TIBBETTS CREEK	8 - Cedar-Sammamish		70112
(View	13138	17110012000221	Water	Bacteria	4A	TIBBETTS CREEK	8 - Cedar-Sammamish	Issaquah Creek Basin Bacteria TMDL	13138
[View]	15779	17110012000224	Water	Bacteria	4.0	TIBBETTS CREEK	8 - Cedar-Sammamish	Issaquah Creek Basin Bacteria THDL	15779
View	13654	17110012000221	Water	Hercury	-2	TIBBETTS CREEK	8 - Cedar Sammamish		13654
View	13645	17110012000221	Water	Ammonia-N	1	TIBBETTS CREEK	8 - Cedar-Sammamish		13645
(View)	13646	17110012000221	Water	Arsenic	1	TIBBETTS CREEK	8 - Cedar-Sammamish		13646
[View]	13657	17110012000221	Water	Selenium	-1	TIBBETTS CREEK	8 - Cedar-Sammamish		13657

303(d) Screen shot

Newport Way Improvements Project Issaquah, Washington



Water Quality Improvement Projects (TMDLs)

Water Quality Improvement > Water Quality Improvement Projects by WRIA 8: Cedar-Sammanish

WRIA 8: Cedar-Sammamish

The following table lists overview information for water quality improvement projects (including total maximum daily loads, or TMDLs) for this water resource inventory area (WRIA). Please use links (where available) for more information on a project.

Counties

- King
- Snohumish



Waterbody Name	Pollutants	Status**	TMDL Lead	
Ballinger Lakë	Total Phosphorus	Approved by EPA	Tricia Shoblom 425-649-7288	
Bear-Evans Creek Basin	Fecal Coliform	Approved by EPA	Joan Nolan	
	Dissolved Oxygen Temperature	Approved by EPA	425-649-4425	
Cottage Lake	Total Phosphorus	Approved by EPA Has an implementation plan	Tricia Shoblom 425-649-7288	
Issaquah Creek Basin	Fecal Coliform	Approved by EPA	30an Nolan 425-649-4425	
Little Bear Creek Tributaries: Trout Stream Great Dane Creek Cutthroat Creek	Fecal Coliform	Approved by EPA	Ralph Svricek 425-649-7036	
North Creek	Fecal Coliform	Approved by EPA Has an implementation plan	Ralph Svricek 425-649-7036	
Pipers Creek	Fecal Coliform	Approved by EPA	Joan Nolan 425-649-4425	
Sammamish River	Dissolved Oxygen Temperature	Field work starts summer 2015	Ralph Svricek 425-649-7036	
Swamp Creek	Fecal Coliform	Approved by EPA Has an implementation plan	Ralph Svricek 425-649-7036	

 $Accessed\ at:\ http://web.archive.org/web/20170330212555/http://www.ecy.wa.gov:80/programs/wq/tmdl/TMDLsbyWria/tmdl-wria08.html$

TMDL Screen Shot

Newport Way Improvements Project Issaquah, Washington



APPENDIX EWetland and Stream Descriptions

TABLE E-1. WETLAND A

Wetland A - Information			
Location	Approximately 30 feet east of Newport Way		
WRIA	8 - Cedar-Sammamish		
Local Jurisdiction	King County	全国国门市	
Category/Rating	II (22 points) ¹		
Buffer Width	100 feet ²	以及其中人名英 克斯	
Size	9.8 acres (estimated)		
Cowardin Class (Cowardin et al. 1979)	Palustrine Forested		
HGM Class	Riverine	2000年1000年100日	
Data Forms	Appendix C: SPs 1-6	Looking northwest towards Wetland A; Newport Way above	
Description Summary			
Vegetation	felix-femina), slough sedge (Carex obnuskunk cabbage (Lysichiton americanus)	is arundinacea), common ladyfern (Athyrium pta), field horsetail (Equisetum arvense) and), willows (Salix spp.), Himalayan blackberry hiphoricarpos albus)	
Soils	Typically meets criteria for redox dark su	rface (F6) and/or loamy mucky mineral (F1).	
Hydrology	facultative (FAC)-neutral test.	igh groundwater table, geomorphic position, ion, high water table, overbank flooding from	
Notes	Riparian wetland on downslope of Newport Way adjacent to Tibbetts Creek.		
Western Washington Wet	land Rating Functions Summary (Append	lix D – 22 points total)	
Water Quality	7 points: due to size, vegetation characte the road (i.e. source of pollution).	eristics and coverage and position adjacent to	
Hydrologic	9 points: due to ground position and ab to a stream that has flooding problems.	ility to detain water that would otherwise flow	
Habitat	6 points: due to numerous vegetation communities, interspersion and habitat features		
Buffer Condition	buffer to the south and west of the wetlar	e traffic volume, is located within the wetland and. The buffer to the east and north is relatively and deciduous forest with a moderately dense er.	

Notes:

- 1. Wetland rating in accordance with Washington State Wetlands Rating System for Western Washington, (Hruby 2014).
- 2. IMC 18.10.640 based on characteristics of a forested wetland and a habitat rating score of 22 points. The final buffer width is subject to approval by the jurisdictional authority.



TABLE E-2. TIBBETTS CREEK

General Information	
Extent of Field Survey	This stream is located at the intersection of Newport Way approximately 200 feet west of SR 900. We characterized the upstream and downstream portions of the stream within the Area of Investigation.
WRIA	8 - Cedar-Sammamish
DNR Stream Type	F
IMC Stream Type	Class 2 Stream with Salmonids
Buffer Width	100 feet
Average Bankfull Width	20 feet
Duration	Perennial
Literature Reviewed	None
Description Summary	
Documented Fish Use	Winter steelhead, kokanee, coho, sockeye
Connectivity	Tibbetts Creek flows into Lake Sammamish, which flows into Lake Washington and ultimately connects to Puget Sound.
Channel Description	Channel consists of gravels and cobbles with sandy silt sediments.
Riparian/Buffer Condition	The buffer has been modified by development on adjacent properties for residential and business uses. Within the survey area, the buffer consists of narrow bands of forested and shrub vegetation. The forested areas are dominated red alder and bigleaf maple. The shrub and understory areas are typically dominated by Himalayan blackberry, with some salmonberry and reed canarygrass.
Existing culvert under Newport Way	There is an existing large box culvert at the crossing of Tibbetts under Newport Way.
Stormwater inputs	There were observed direct discharge points of stormwater into Tibbetts Creek.
Notes	Within this section of Tibbetts Creek, habitat and water quality functions have been degraded with invasive species dominant in the riparian buffer.



TABLE E-3. STREAM 0169G

General Information		
Extent of Field Survey	This stream is located approximately 500 feet west of the intersection of Newport Way and SR 900. We characterized the upstream and downstream portions of the stream within the Area of Investigation as well as extending approximately 650 feet upstream of the Newport Way crossing.	
WRIA	8 - Cedar-Sammamish	
Local Jurisdiction	King County	
DNR Stream Type	NA (not mapped)	
Local Jurisdiction Stream Type	Class 2 Stream with Salmonids	
Buffer Width	100 feet	
Average Bankfull Width	4 feet	
Duration	Seasonal	
Reference Studies	None	
Description Summary		
Documented Fish Use	None	
Connectivity	This tributary connects with Tibbets Creek. The existing road crossing is a fish passage barrier.	
Channel Description	Downstream channel adjacent to the culvert is predominantly vegetated with grass and very limited substrate evident. Further downstream, the channel has been modified, including installation of pavers to harden the stream bank. Upstream portion of the channel was not clearly observed from ROW.	
Riparian/Buffer Condition	The buffer is severely degraded and is extensively modified by the adjacent residence, including associated lawn areas. A large amount of debris and litter was observed throughout the buffer. The buffer consists of narrow bands of forested and shrub vegetation, with some areas that have been cleared and are now limited to grass and other herbaceous species. The forested areas are dominated by black cottonwood with some red alder and Pacific willow (Salix lasiandra). The shrub and understory areas are typically dominated by Himalayan blackberry and reed canarygrass.	
Existing culvert under Newport Way	According to a 2017 Culvert Inventory (City of Issaquah 2017b) provided by City of Issaquah Public Works, this stream is conveyed under Newport Way through an 18-inch pipe culvert.	
Stormwater inputs	Based on field observations, stormwater discharges untreated from the roadway directly into the downstream portion of this tributary.	
Notes	The western portion of this stream, on the other side of Newport Way, was not delineated due to ROE limitations. This portion of the stream is within a predominantly undisturbed forested hillside.	



TABLE E-4. STREAM 0169H

General Information		
Extent of Field Survey	This stream is located approximately 0.45 miles west of the intersection of Newport Way and SR 900 on King County owned property. The upstream portion of this property was characterized in the field. Observations of the downstream portion were limited to the ROW.	
WRIA	8 - Cedar-Sammamish	
DNR Stream Type	NA (not mapped)	
IMC Stream Type	Class 2 Stream with Salmonids	
Buffer Width	100 feet	
Average Bankfull Width	8-9 feet	
Duration	Seasonal	
Reference Studies	None	
Description Summary		
Documented Fish Use	None	
Connectivity	The stream flows through a box culvert under Newport Way before connecting downstream with Tibbetts Creek. The culvert is a fish barrier.	
Channel Description	Channel consists of large boulders, cobbles and gravels with very minor sandy silt sediments.	
Riparian/Buffer Condition	This buffer is predominantly undisturbed with some elements of degradation with invasive species and litter from the roadway. The buffer consists of forested and shrub vegetation, with some areas that have been cleared and are now limited to grass and other herbaceous species. The forested areas are dominated by black cottonwood with some red alder and Pacific willow. The shrub and understory areas are typically dominated by Himalayan blackberry, with some snowberry (<i>Symphoricarpos albus</i>), salmonberry, reed canarygrass, English ivy (<i>Hedera helix</i>) and sword fern (<i>Polystichum munitum</i>).	
Existing culvert under Newport Way	According to a 2017 Culvert Inventory (City of Issaquah 2017b) provided by City of Issaquah Public Works, this stream flows through a 36-inch reinforced concrete pipe under Newport Way.	
Stormwater Inputs	Currently, stormwater flows untreated from the roadway into the stream.	
Notes	Due to ROE constraints, the survey was limited to the upstream portion of the tributary. This tributary is located on King County parcel and part of the Cougar Mountain Regional Wildland Park.	



TABLE E-5. STREAM 0169I

General Information	
Extent of Field Survey	This stream is located approximately 0.25 miles west of the intersection of Newport Way and SR 900. We characterized the upstream and downstream portions of the stream within the Area of Investigation as well as extending approximately 650 feet upstream of the Newport Way crossing .
WRIA	8 - Cedar-Sammamish
DNR Stream Type	NA (not mapped)
IMC Stream Type	Class 3
Buffer Width	50 feet
Average Bankfull Width	2-3 feet
Duration	Seasonal
Reference Studies	Bergsma Plan Set (Pace 2016), Stream 2 Survey (CEC 2016)
Description Summary	
Documented Fish Use	None
Connectivity	There is a pipe culvert under Newport Way; the downstream channel flows a short distance before entering Wetland A, through which there is no contiguous stream channel.
Channel Description	Downstream channel section enters wetland approximately 35 feet downstream of the culvert crossing. There is no contiguous channel through the wetland. Substrate is sand and gravel. Upstream channel section is steep and narrow, with sand and gravel dominant.
Riparian/Buffer Condition	This buffer is predominantly undisturbed, and vegetation is predominantly forested and shrub. The shrub and understory areas are typically dominated by Himalayan blackberry, with some snowberry (Symphoricarpos albus) and sword fern (Polystichum munitum).
Existing Culvert under Newport Way	According to a 2017 Culvert Inventory (City of Issaquah 2017b) provided by City of Issaquah Public Works, this culvert is 18-inch reinforced concrete pipe.
Stormwater Inputs	Currently, stormwater flows untreated from the roadway into the stream.
Notes	Survey was limited to the downstream portion of the tributary; upstream OHWM obtained from developer drawings.



TABLE E-6. ANTI-AIRCRAFT CREEK

General Information	
Extent of Field Survey	Anti-Aircraft is located approximately 200 feet south of intersection of Newport Way and NW Oakcrest Drive. Neither the upstream nor downstream portions of Anti-Aircraft Creek were included in our survey due to recent re-construction of the stream channel, resulting in absence of OHWM indicators.
WRIA	8 - Cedar-Sammamish
DNR Stream Type	F
IMC Stream Type	Class 2 Stream with Salmonids
Buffer Width	100 feet
Average Bankfull Width	Not characterized
Duration	Seasonal
Reference Studies	Culvert Replacement Project Design Drawings (PBS 2017)
Description Summary	
Documented Fish Use	Fall Chinook, coho, sockeye, winter steelhead
Connectivity	Connects with Tibbetts Creek downstream of Newport Way.
Riparian Buffer Condition	Vegetation has been disturbed and replaced as part of ongoing restoration activities.
Existing culvert under Newport Way	Based on 2017 Newport Culvert Inventory (City of Issaquah 2017b) provided by City of Issaquah Public Works, a 20-foot box culvert has been constructed to replace the prior 36-inch reinforced concrete pipe crossing.
Stormwater Inputs	Not assessed
Notes	This stream has undergoing recent restoration, including re-routing the channel and a new culvert crossing under Newport Way.



TABLE E-7. SCHNEIDER CREEK

General Information		
Extent of Survey	This stream is located approximately 0.84 miles west along Newport Way from the intersection of Newport Way and SR 900. Our field survey included both upstream and downstream portions within the Area of Investigation.	
WRIA	8 - Cedar-Sammamish	
DNR Stream Type	F	
IMC Stream Type	Class 2 Stream with Salmonids	
Buffer Width	100 feet	
Average Bankfull Width	10 feet	
Duration	Perennial	
Reference Studies	Gateway Apartments Critical Areas Study (Talasaea 2014)	
Description Summary		
Documented Fish Use	Cutthroat	
Connectivity	Schneider Creek flows under Newport Way via a 3-foot round culvert and then ultimately into Lake Sammamish. The existing crossing is a total fish barrier.	
Channel Description	Upstream substrate consists of cobbles, gravel and sand. Downstream channel contains some large cobbles at the culvert outlet and then transitions to gravels and cobbles.	
Riparian/Buffer Condition	Above the culvert, the stream has a well-established forested riparian buffer with predominantly native species. There is an increase in invasive species within the riparian buffer as the stream gets closer to Newport Way. The forested areas are dominated by black cottonwood with some red alder and Pacific willow (Salix lasiandra). The shrub and understory areas are typically dominated by Himalayan blackberry, with some snowberry, salmonberry, reed canarygrass and sword fern. Below the culvert at Newport Way, the stream opens to a highly degraded riparian buffer adjacent to a residential property with lawn growing up to the bank of the stream. Beyond this property, the stream flows into a forested area dominated by black cottonwood with some red alder and Pacific willow. The shrub and understory areas are typically dominated by Himalayan blackberry, with some snowberry, salmonberry (Rubus spectabilis), reed canarygrass (Phalaris arundinacea) and sword fern (Polystichum munitum).	
Existing Culvert under Newport Way	According to a 2017 Culvert Inventory (City of Issaquah 2017b) provided by City of Issaquah Public Works, this culvert is a 30-inch corrugated metal pipe.	
Stormwater Inputs	Currently, stormwater flows untreated from the roadway into the channel.	
Notes	Within this section of the stream, habitat and water quality functions have been degraded. There is little to no vegetated buffer downstream of Newport Way, and human use and alteration are evident.	



